

ADA 077832



9 14 CNA-PP-7
PROFESSIONAL PAPER 258 November 1979

11/78 - Jan 79

12 94

6
**ANALYTICAL METHODS IN
SEARCH THEORY.**

10
Marc S. Mangel
James A. Thomas, Jr.



CENTER FOR NAVAL ANALYSES

2000 North Beauregard Street, Alexandria, Virginia 22311

Reproduced From
Best Available Copy

79 127045
270 100

B

PROFESSIONAL PAPER 258 / November 1979

ANALYTICAL METHODS IN SEARCH THEORY

Marc S. Mangel
James A. Thomas, Jr.



Operations Evaluation Group

CENTER FOR NAVAL ANALYSES

2000 North Beauregard Street, Alexandria, Virginia 22311

| | |
|------------------|-------------------------|
| Distribution | |
| None | |
| DCC TAB | |
| Unclassified | |
| Justification | |
| By | |
| Distribution | |
| Additional Codes | |
| DIST. | Available/or Special |

A handwritten signature or mark is present in the bottom left corner of the form.

PREFACE

This work is an expanded version of a set of lecture notes corresponding to a course given by the author during the period November 1978 to January 1979. Analytical methods for the solution of moving object search problems are developed from "first principles." This work complements the first author's CNA memorandum, (reference 1) ✓ but otherwise there is little overlap with the existing literature on search theory (e.g., references 2, 3, and 4). The approach taken is tutorial, in that the solutions of harder problems are motivated by the solutions of simpler problems. There are exercises interspersed throughout the work; the solutions of some of these are given in the appendix.

This work is concerned with mathematical analysis and not with modeling target motion or detection functions and not with the development of computer codes.

Some general references are references 2 through 7.

TABLE OF CONTENTS

| | <u>Page</u> |
|--|-------------|
| Formulation of the Search Problem | 1 |
| Initial Density | 1 |
| Type 1: Initial Density Concentrated at a Point. | 2 |
| Type 2: Nowhere Vanishing Initial Densities. | 3 |
| Type 3: Compact Initial Data | 3 |
| Target Motion Model | 3 |
| Deterministic Target Motion | 5 |
| Conditionally Deterministic Target Motion | 6 |
| Stochastic Target Motion. | 6 |
| Search Function | 8 |
| Search Problems | 9 |
| The Search Equations. | 10 |
| Scaled Search Equation. | 14 |
| Solution of the Deterministic and Conditionally Deterministic Search Equations. | 18 |
| Computational Algorithm for Deterministic Target Motion | 22 |
| Computational Algorithm for Conditionally Deterministic Target Motion | 22 |
| Alternative Computational Algorithm | 24 |
| Solution of the Stochastic Search Equation in Some Special Cases | 25 |
| Canonical Problem: Type 1 Initial Data | 26 |
| Canonical Problem: General Initial Data. | 27 |
| Canonical Problem: Type 3 Initial Data | 28 |
| Canonical Problem: Nonconstant ψ | 29 |
| Canonical Problem: Ornstein-Uhlenbeck Process. | 30 |
| Approximate Solution of the Stochastic Search Equation in the General Case. | 31 |
| Alternative Algorithm for the Solution of the Hamilton-Jacobi Equation. | 35 |
| Algorithm for Solution of the Hamilton-Jacobi Equation. | 36 |
| Variational Interpretation. | 37 |
| Initial Condition | 41 |
| Summary of Computational Algorithms | 42 |
| Deterministic Target Motion | 42 |
| Conditionally Deterministic Target Motion | 43 |
| Stochastic Target Motion. | 43 |

TABLE OF CONTENTS
(Continued)

| | <u>Page</u> |
|--|-------------|
| Optimal Search for a Moving Target. | 45 |
| Functional #1: Instantaneous Probability of Detection. | 45 |
| Functional #2: Probability of Detection by Time T_S | 46 |
| Functional #3: Mean Time to Detection. | 47 |
| Solution of an Optimal Search Problem | 47 |
| References. | 50 |
| Appendix A: Solution of Exercises | |

FORMULATION OF THE SEARCH PROBLEM

The position of the target at time t will be denoted by $x(t)$. It is a vector, so that:

$$x(t) = (x_1(t), x_2(t), x_3(t)) . \quad (1.1)$$

The target moves in some region, denoted by D_T .

The position of the searcher at time t will be denoted by:

$$z(t) = (z_1(t), z_2(t), z_3(t)) . \quad (1.2)$$

The following inputs are needed to formulate the search problem.

INITIAL DENSITY

Let $\rho_0(x)$ be defined by:

$$\begin{aligned} \rho_0(x)dx &= \Pr\{x_1 \leq X_1(0) \leq x_1 + dx_1, x_2 \leq X_2(0) \leq x_2 + dx_2, \\ &\quad x_3 \leq X_3(0) \leq x_3 + dx_3\} . \end{aligned} \quad (1.3)$$

In this equation, $x = (x_1, x_2, x_3)$ is a vector and $dx = (dx_1, dx_2, dx_3)$. Three types of initial densities are considered.

Type 1

Initial density concentrated at a point. Namely:

$$\text{Prob}\{X(0) = x_0\} = 1. \quad (1-4)$$

Symbolically, one writes:

$$\rho_0(x) = \delta(x-x_0) = \delta(x_1-x_{10})\delta(x_2-x_{20})\delta(x_3-x_{30}). \quad (1-5)$$

In this equation, $\delta(s)$ is the "Dirac Delta Function" (reference 8).

Exercise (See Reference 8)

One definition of $\delta(s)$ is:

$$\delta(s) = \lim_{n \rightarrow \infty} \sqrt{\frac{n}{2\pi}} e^{-s^2 n/2} \quad (1-6)$$

$$= \lim_{n \rightarrow \infty} \epsilon_n(s). \quad (1-7)$$

Sketch a few of the $\epsilon_n(s)$. Also draw $\delta'(s)$, the derivative of $\delta(s)$.

Note that $\delta(s) = 0$ if $s \neq 0$ and:

$$\int \delta(s) ds = 1. \quad (1-8)$$

$$\int h(s) \delta(s) ds = h(0). \quad (1-9)$$

Type 2: Nowhere Vanishing Initial Densities

In this case, $\rho_0(x)$ does not vanish in D_T . An example of such an initial density is the circular Gaussian:

$$\rho_0(x) = \left[\frac{1}{2\pi\sigma^2} \right]^{3/2} e^{-r^2/2\sigma^2}, \quad (1-10)$$

where $r = (x_1^2 + x_2^2 + x_3^2)^{\frac{1}{2}}$.

Type 3: Compact Initial Data

In this case, $\rho_0(x)$ is non-zero in some subregion $D_T' \subseteq D_T$ and zero elsewhere (see figure 1).

TARGET MOTION MODEL

The velocity of the target is $v(x,t)$ and is given by:

$$\frac{dx}{dt} = v(x,t). \quad (1-11)$$

The increment in x is then given by $\Delta x = v(x,t)\Delta t$. Introduce the transition function (e.g., references 5 and 6), $q(\xi, t, \Delta t, x)$ defined as follows.

Set:

$$\Delta x = x(t+\Delta t) - x(t). \quad (1-12)$$

Then, the transition function $q(\xi, t, \Delta t, x)$ is:

$$q(\xi, t, \Delta t, x) d\xi = \text{Prob}\{\xi \leq \Delta x \leq \xi + d\xi \mid X(t) = x\}. \quad (1-13)$$




FIG. 1: ILLUSTRATION OF D_T AND TYPE 3 INITIAL DATA

In this equation, $\xi = (\xi_1, \xi_2, \xi_3)$ and $d\xi = (d\xi_1, d\xi_2, d\xi_3)$.

The right-hand side of equation 1-13 is satisfied by each component.

Equation 1-13 can also be rewritten as:

$$q(\xi, t, \Delta t, x) d\xi = \text{Prob}\{\xi \leq X(t+\Delta t) - X(t) \leq \xi + d\xi | X(t) = x\}. \quad (1-14)$$

Deterministic Target Motion

In the case of deterministic target motion,

$$q(\xi, t, \Delta t, x) = \delta(\xi - b(x, t)\Delta t). \quad (1-15)$$

Namely, with probability 1, $\Delta X = b(x, t)\Delta t$. Let $E\{\cdot\}$ denote mathematical expectation. Then:

$$\begin{aligned} E(\Delta X | X(t) = x) &= \int \xi q(\xi, t, \Delta t, x) d\xi \\ &= \int \xi \delta(\xi - b(x, t)\Delta t) d\xi \\ &= b(x, t)\Delta t. \end{aligned} \quad (1-16)$$

Hence, one writes that:

$$\frac{dx}{dt} = b(x, t). \quad (1-17)$$

Exercise

Show that if $q(t, t', x)$ is given by equation 1-16, then

$$\text{Var}(X|X(t) = x) = 0.$$

Conditionally Deterministic Target Motion

In this case, equation 1-15 is replaced by:

$$q(t, t, \Delta t, x) = \sum_{\alpha} p_{\alpha} \delta(t - b^{\alpha}(x, t)\Delta t), \quad (1-18)$$

with $\sum_{\alpha} p_{\alpha} < 1$. Namely, there are a number of possible target velocities, with a probability p_{α} associated with velocity $b^{\alpha}(x, t)$.

If the velocities are continuously distributed, then equation 1-18 must be modified, of course.

The classical example of conditionally deterministic target motion is the "fleeing datum" (see reference 2). In this case, the target flees with a known velocity but with unknown bearing.

Stochastic Target Motion

In this case, one assumes that given $x(t) = x$, then $\Delta X = X(t+\Delta t) - X(t)$ is normally distributed with mean $\hat{b}(x, t)\Delta t + o(\Delta t)$ *

*Without loss of generality, one assumes $\hat{b}(x, t)$ is known with complete certainty. Only a small modification is needed to use a conditionally deterministic mean $\hat{b}^{\alpha}(x, t)$.

and covariance matrix $\hat{a}_{ij}^{\Delta t}(x, t)\Delta t + o(\Delta t)$. Here $o(\Delta t)$ means a quantity such that:

$$\lim_{\Delta t \rightarrow 0} \frac{o(\Delta t)}{\Delta t} = 0. \quad (1-19)$$

Alternatively, one writes that:

$$E(\Delta X_i | X(t) = x) = \hat{b}_i(x, t)\Delta t + o(\Delta t) \quad i = 1, 2, 3 \quad (1-20)$$

$$\text{Cov}(\Delta X_i \Delta X_j | X(t) = x) = \hat{a}_{ij}^{\Delta t}(x, t)\Delta t + o(\Delta t). \quad (1-21)$$

Exercise

Show that:

$$E(\Delta X_i \Delta X_j | X(t) = x) = \hat{a}_{ij}^{\Delta t}(x, t)\Delta t + o(\Delta t) \quad (1-22)$$

Exercise

Let $\Delta t \rightarrow 0$ and then $(\hat{a}_{ij}^{\Delta t}) \rightarrow 0$. What happens to the density $q(\xi, t, \Delta t, x)$ for ΔX ?

Exercise

Restate equations 1-20 and 1-22 in terms of $q(\xi, t, \Delta t, x)$.

Remark 1

1. Since for stochastic target motion

$$d\hat{x} \sim N(b(x,t)\Delta t + o(\Delta t), a(x,t)\Delta t + o(\Delta t)) , \quad \hat{x}(t)$$

satisfies the "Ito Equation" $d\hat{x} = b(x,t)dt + \sqrt{a(x,t)} dW$,
where $W(t)$ is Brownian motion.

2. For the case of stochastic target motion

$$\int \hat{x}_n^q(r,t,\Delta t,x) dr = o(\Delta t) \quad \text{for } n \geq 3 .$$

Exercise

Demonstrate Remark 2.

SEARCH FUNCTION

The final input is the search function (or "conditional detection function", reference 9). Let:

$$\hat{\psi}(x,t,z)\Delta t = \text{Prob}\{\text{detection in } (t, t+\Delta t) | X(t)=x, Z(t)=z\}. \quad (1-23)$$

This function must be modeled.

SEARCH PROBLEMS

The following quantities are of interest in search theory.

Let:

$$\rho(x, t | z) dx = \Pr\{x \leq X(t) \leq x + dx | \text{search along } Z(\tau), \\ 0 \leq \tau \leq t \text{ was not successful}\} . \quad (1-24)$$

To obtain $\rho(x, t | z)$, consider the joint density

$$f(x, t; z) dx = \Pr\{x \leq X(t) \leq x + dx \text{ and search along } Z(\tau), \\ 0 \leq \tau \leq t \text{ was not successful}\} . \quad (1-25)$$

Since $\Pr(A \text{ and } B) = \Pr(A|B)\Pr(B)$, one has:

$$\rho(x, t | z) = \frac{f(x, t; z)}{\int_{D_T} f(x, t; z) dx} . \quad (1-26)$$

Note that $\rho(x, t | z)$ and $f(x, t; z)$ have the same spatial dependence and differ only by a function of time.

A related quantity, of interest in "time-late" problems, is the density in the absence of search:

$$\rho(x, t) dx = \Pr\{x \leq X(t) \leq x + dx\} . \quad (1-27)$$

THE SEARCH EQUATIONS

General References: 10, 11, 12.

In order to derive the search equation, consider the time increment $(t, t + \Delta t)$. Then, in this increment, one finds that

$$f(x, t + \Delta t; z) dx = \int \left(1 - \hat{\psi}(x - \xi, t; z) \Delta t \right) q(\xi, t, \Delta t, x - \xi) f(x - \xi, t; z) d\xi dx. \quad (2-1)$$

When the right-hand side is Taylor expanded, one obtains:

$$\begin{aligned} f(x, t + \Delta t; z) dx &= \int \left\{ \left(1 - \hat{\psi}(x, t; z) \Delta t \right) \left[q(\xi, t, \Delta t, x) f(x, t; z) \right. \right. \\ &\quad \left. - \sum_i \xi_i \frac{\partial}{\partial x_i} (qf) + \frac{1}{2} \sum_{ij} \xi_i \xi_j \frac{\partial^2}{\partial x_i \partial x_j} (qf) \right] \\ &\quad \left. + \Delta t q(\xi, t, \Delta t, x - \xi) f(x - \xi, t; z) \cdot O(\xi) + O(\xi^3) \right\} d\xi dx \end{aligned} \quad (2-2)$$

$$\begin{aligned} &= (1 - \hat{\psi} \Delta t) \left\{ \int d\xi q(\xi, t, \Delta t, x) f(x, t; z) \right. \\ &\quad \left. - \sum_i \frac{\partial}{\partial x_i} (\xi_i qf) + \sum_{ij} \frac{1}{2} \frac{\partial^2}{\partial x_i \partial x_j} (\xi_i \xi_j qf) + O(\xi^3) \right\} \quad (2-3) \\ &\quad + \Delta t \int q(\xi, t, \Delta t, x - \xi) f(x - \xi, t; z) O(\xi) d\xi dx . \end{aligned}$$

$$\begin{aligned}
 f(x, t+\Delta t; z) &= (1 - \hat{\psi} \Delta t) \left[f(x, t; z) \int q(\xi, t, \Delta t, x) d\xi \right. \\
 &\quad - \sum_i \frac{\partial}{\partial x_i} \left(f \int \xi_i q(\xi, t, \Delta t, x) d\xi \right) \\
 &\quad + \sum_{ij} \frac{1}{2} \frac{\partial^2}{\partial x_i \partial x_j} \left(f \int \xi_i \xi_j q(\xi, t, \Delta t, x) d\xi \right) \\
 &\quad \left. + \int O(\xi^3) q(\xi, t, \Delta t, x) d\xi \right] + o(\Delta t) .
 \end{aligned} \tag{2-4}$$

Thus:

$$\begin{aligned}
 f(x, t+\Delta t; z) &= (1 - \hat{\psi} \Delta t) \left[f(x, t; z) - \sum_i \frac{\partial}{\partial x_i} \left(\hat{b}_i(x, t) \Delta t + o(\Delta t) \right) \right. \\
 &\quad \left. + \frac{1}{2} \sum_{ij} \frac{\partial^2}{\partial x_i \partial x_j} \left(\hat{a}_{ij}(x, t) \Delta t + o(\Delta t) \right) + o(\Delta t) \right] .
 \end{aligned} \tag{2-5}$$

Equation 2-5 follows from the definition of $\hat{b}(x, t)$ and $\hat{a}(x, t)$.

Expanding the right-hand side gives:

$$\begin{aligned}
 f(x, t+\Delta t; z) - f(x, t; z) &= -\hat{\psi} f \Delta t - \sum_i \frac{\partial}{\partial x_i} (\hat{b}_i f) \Delta t \\
 &\quad + \frac{1}{2} \sum_{ij} \frac{\partial^2}{\partial x_i \partial x_j} (\hat{a}_{ij} f) \Delta t + o(\Delta t) .
 \end{aligned}$$

Dividing by Δt and taking the limit as $\Delta t \rightarrow 0$ gives the search equation:

$$\frac{\partial f}{\partial t} = \frac{1}{2} \sum_{i,j} \frac{\partial^2}{\partial x_i \partial x_j} (\hat{a}_{ij} f) - \sum_i \frac{\partial}{\partial x_i} (\hat{b}_i f) - \hat{\psi} f . \quad (2-7)$$

Equation 2-7 will be called the stochastic search equation (SSE).

Exercise

Suppose that there is no search. Let

$\rho(x, t) dx = \Pr\{x \leq X(t) \leq x + dx\}$. Show that $\rho(x, t)$ satisfies the following equation:

$$\frac{\partial \rho}{\partial t} = \sum_{i,j} \frac{1}{2} \frac{\partial^2}{\partial x_i \partial x_j} (\hat{a}_{ij} f) - \sum_i \frac{\partial}{\partial x_i} (\hat{b}_i f) . \quad (2-8)$$

Exercise

For the case of deterministic target motion, show that

$f(x, t; z)$ satisfies the deterministic search equation (DSE):

$$\frac{\partial f}{\partial t} = - \sum_i \frac{\partial}{\partial x_i} (\hat{b}_i f) - \hat{\psi} f . \quad (2-9)$$

Remark

Define a flux $J_i(x, t; f)$ by:

$$J_i(x, t; f) = - \left[\sum_j \frac{\partial}{\partial x_j} (\hat{a}_{ij} f) - \hat{b}_i f \right]. \quad (2-10)$$

Then, the SSE (equation 2-7) becomes:

$$\frac{\partial f}{\partial t} + \sum_{i=1}^n \frac{\partial}{\partial x_i} [J_i(x, t; f)] = - \hat{\psi} f. \quad (2-11)$$

This is a conservation or continuity equation.

Now consider equation 2-7. It needs boundary and initial conditions.

1. Initial Condition: $f(x, t; z) \rightarrow p_0(x)$ as $t \rightarrow 0$,
where $p_0(x)$ is the initial density.
2. Boundary Conditions are more difficult. Possible ones
are:
 - a. $f(x, t; z) = 0$ for x on the boundary of D_T
(i.e., an absorbing boundary).
 - b. The normal derivative $\partial f / \partial n = 0$ on the boundary
(reflecting boundary).

In this paper, the boundaries are simply ignored. This
leads to solutions that are valid far away from the boundary.

They can be modified to satisfy any needed boundary conditions
(see reference 13 or 1).

Remark

$f(x, t; z)$ does not integrate to 1.

In fact,

$$\int_{D_T} f(x, t; z) dx = \text{Prob}\{\text{search up to time } t \text{ is not successful}\} + 1 \text{ (hopefully)}.$$

THE SCALED SEARCH EQUATION

Let T_c , L_c , b_m , a_m be defined as follows:

T_c = a characteristic time, e.g., time available for search

L_c = a characteristic distance, e.g., distance from
center of $\hat{v}_0(x)$ to the boundary

a_m = maximum value of $\hat{a}_{ij}(x, t)$ over all i, j

b_m = maximum value of $\hat{b}_i(x, t)$ over all i, j .

Now define dimensionless variables by:

$$\tau = \frac{t}{T_c} \quad y_i = \frac{x_i}{L_c} \quad \hat{a}_{ij} = \frac{\hat{a}_{ij}}{a_m} \quad \hat{b}_i = \frac{\hat{b}_i}{b_m}$$

The search equation 2-7 takes the form:

$$\frac{1}{T_c} \frac{\partial f}{\partial \tau} = \frac{a_m^T}{L_c^2} \sum_{i,j} \frac{\partial^2}{\partial y_i \partial y_j} (a_{ij} f) - \frac{b_m^T}{L_c} \sum_i \frac{\partial}{\partial y_i} (\bar{b}_i f) - \hat{\psi} f , \quad (2-12)$$

or

$$\frac{\partial f}{\partial \tau} = \frac{a_m^T}{L_c^2} \sum_{i,j} \frac{\partial^2}{\partial y_i \partial y_j} (a_{ij} f) - \frac{b_m^T}{L_c} \sum_i \frac{\partial}{\partial y_i} (\bar{b}_i f) - \hat{\psi} T_c f . \quad (2-13)$$

Now assume that T_c, L_c are chosen so that:

$$\frac{a_m^T}{L_c^2} \equiv \epsilon \ll 1 . \quad (2-14)$$

Define:

$$\left. \begin{array}{l} b_i = \bar{b}_i \left(\frac{b_m^T}{L_c} \right) \\ \psi = \hat{\psi} T_c \end{array} \right\} \quad (2-15)$$

The final non-dimensional search equations is (reverting to x, t instead of y, τ):

$$\frac{\partial f}{\partial t} = \epsilon \sum_{i,j} \frac{\partial^2}{\partial x_i \partial x_j} (a_{ij} f) - \sum_i \frac{\partial}{\partial x_i} (b_i f) - \psi f . \quad (2-16)$$

Remark

When $\epsilon \rightarrow 0$, equation 2-16 becomes the deterministic search equation.

Remark

It has been assumed that b_i and ψ are quantities that are $O(1)$, i.e., of the order of 1. It could be that b_i are $O(\epsilon)$, in which case equation 2-16 would take the form:

$$\frac{\partial f}{\partial t} = \epsilon \left[\sum_{i,j} \frac{\partial^2}{\partial x_i \partial x_j} (a_{ij} f) - \sum_i \frac{\partial}{\partial x_i} (b_i f) \right] - \psi f . \quad (2-17)$$

For purposes here, equation 2-16 will be used instead of equation 2-17.

Remark: An Alternative Formulation for ϵ

Suppose that the initial data were:

$$p_0 = \frac{1}{2\pi\sigma^2} \exp \left[- \frac{(x_1^2 + x_2^2)}{2\sigma^2} \right] .$$

Under the scaling

$$x_i = y_i L_c ,$$

one obtains:

$$p_0 = \frac{1}{2\pi\sigma^2} \exp \left[- \frac{(y_1^2 + y_2^2) L_c^2}{2\sigma^2} \right] .$$

If $\sigma^2/2L_c^2 \ll 1$, then an alternative definition of ϵ could be:

$$\epsilon = \frac{\sigma^2}{2L_c^2}$$

In this case, (a_{ij}) and $b(y,t)$ would have different interpretations.

Remark

One could imagine that $b(x,t)$ in equation 2-16 is not known with complete certainty, but that $b(x,t)$ is known in a conditionally deterministic fashion. In this case, one solves an equation of the form of equation 2-16 for each possible $b(x,t)$ and then averages over a .

SOLUTION OF THE DETERMINISTIC AND
CONDITIONALLY DETERMINISTIC SEARCH EQUATIONS

References: 14,15.

The deterministic search equation is:

$$\frac{\partial f}{\partial t} = - \sum_{i=1}^n \frac{\partial}{\partial x_i} (b_i f) - \psi f , \quad (3-1)$$

with initial data:

$$f(x, 0; z) = p_0(x), \quad (3-2)$$

and no boundary data.

In order to solve equation 3-1, the method of characteristics
is used. Rewrite equation 3-1 as:

$$\frac{\partial f}{\partial t} + \sum b_i \frac{\partial f}{\partial x_i} = - \left(\psi + \sum \frac{\partial b_i}{\partial x_i} \right) f . \quad (3-3)$$

Introduce a new variable s so that:

$$\frac{dt}{ds} = 1 , \quad (3-4)$$

and set:

$$\frac{dx_1}{ds} = b_1(x, t(s)) . \quad (3-5)$$

Choose initial conditions so that:

$$\left. \begin{array}{l} t = 0 \quad \text{when } s = 0 \\ x_i = x_{i0} \quad \text{when } s = 0 \end{array} \right\} . \quad (3-6)$$

The solutions of equations 3-4 and 3-6 give a "curve" (or space time ray) in D_T . From equation 3-3:

$$\frac{df}{ds} = \frac{\partial f}{\partial t} \frac{dt}{ds} + \sum \frac{\partial f}{\partial x_i} \frac{dx_i}{ds} \quad (3-7)$$

$$= \frac{\partial f}{\partial t} + \sum b_i \frac{\partial f}{\partial x_i} . \quad (3-8)$$

Thus, on the solution curves of equations 3-4 and 3-5,

$$\frac{df}{ds} = - \left[\psi(x(s), s, z(s)) + \sum \frac{\partial b_i}{\partial x_i} (x(s), s) \right] f \quad (3-9)$$

with

$$f = p_0(x_0) \quad \text{when } s = 0 \quad (3-10)$$

(see figure 2).

Formally, one can write:

$$f(x, t; z) = p_0(x_0) \exp \left[- \int_0^t \left(\psi + \sum \frac{\partial b_i}{\partial x_i} \right) ds \right] . \quad (3-11)$$




FIG. 2: SOLUTION OF THE DETERMINISTIC SEARCH EQUATION

In using this formula, one writes:

$$x_i(t) = x_{i0} + \int_0^t b_i(x, s) ds \quad (3-12)$$

or

$$x_{i0} = x_i(t) - \int_0^t b_i(x, s) ds . \quad (3-13)$$

This form is especially useful when b is independent of x . In that case:

$$x_0 = x - \int_0^t b(s) ds . \quad (3-14)$$

EXERCISE (A CASE WHERE EQUATION 3-13 IS USEFUL)

Suppose that $b_i = v_i(t)$, independent of x . Show that:

$$f(x, t; z) = p_0 \left[x - \int_0^t v(s) ds \right] \exp \left\{ - \int_0^t \psi \left[x - \int_s^t v(s') ds' , s, z \right] ds \right\} \quad (3-15)$$

Write down the integral that gives the probability of detection by time t .

Remark

Note that equations 3-11 and 3-14 can be summarized as:

$$f \sim [\text{term due to target motion}] \times [\text{term due to search}] .$$

COMPUTATIONAL ALGORITHM FOR DETERMINISTIC TARGET MOTION

Step 1

Pick $x_0 = (x_{10}, x_{20}, \dots, x_{j0})$

Step 2

Solve $\frac{dx_i}{dt} = b_i(x, t) ; \quad x_i(0) = x_{i0}$

Step 3

Solve $\frac{df}{dt} = -\left(\psi + \sum_i \frac{\partial b_i}{\partial x_i}\right) f$

$$f(0) = \rho_0(x_0).$$

Step 4

Cycle (i.e., pick a different x_0 and return to Step 1).

COMPUTATIONAL ALGORITHM FOR CONDITIONALLY DETERMINISTIC TARGET MOTION

In this case

$$b(x, t) = b^\alpha(x, t) \quad (3-16)$$

with probability p_α . Let $f^\alpha(x, t; z)$ satisfy:

$$\frac{\partial f^\alpha}{\partial t} = -\sum_i \frac{\partial}{\partial x_i} (b_i^\alpha f^\alpha) - \psi f^\alpha \quad (3-17)$$

$$f^\alpha(x, 0; z) = \rho_0(x). \quad (3-18)$$

In order to determine $f(x, t; z) = E_\alpha\{f^\alpha(x, t; z)\}$, the following computational algorithm can be used.

Step 1

Fix $b^\alpha(x, t)$ (with probability p_α).

Step 2

Pick $x_0 = (x_{10}, x_{20}, x_{30})$

Step 3

Solve: $\frac{dx_i}{dt} = b_i^\alpha(x, t); \quad x_i(0) = x_{i0}$.

Step 4

Solve: $\frac{df^\alpha}{dt} = -\left(\psi + \sum_i \frac{\partial b_i^\alpha}{\partial x_i}\right)f^\alpha \quad f^\alpha(0) = p_0(x_0)$.

Step 5

Cycle to step 2 as desired.

Step 6

Cycle through $b^\alpha(x, t)$.

Step 7

Construct: $f(x, t; z) = E_\alpha\{f^\alpha(x, t; z)\}$.

ALTERNATIVE COMPUTATIONAL ALGORITHM

Step 1

Fix: $x_0 = (x_{10}, x_{20}, x_{30})$

Step 2

Fix: $b^\alpha(x, t)$

Step 3

Solve: $\frac{dx_i}{dt} = b_i^\alpha(x, t) \quad x_i(0) = x_{i0}$

Step 4

Solve: $\frac{df^\alpha}{dt} = -\left(\psi + \sum \frac{\partial b_i^\alpha}{\partial x_i}\right) f^\alpha$

Step 5

Cycle through $b^\alpha(x, t)$

Step 6

Construct: $f(x, t; z) = E_\alpha \{f(x, t; z)\}$

Step 7

Cycle through x_0 .

SOLUTION OF THE STOCHASTIC SEARCH EQUATION
IN SOME SPECIAL CASES

General References: 6, 13, 15, 16, 17, 18.

In order to motivate the solutions constructed in the next section, certain special cases are studied in this section. Study of such canonical problems (see reference 16) has been very useful in other disciplines.

First consider the stochastic search equation:

$$\frac{\partial f}{\partial t} = \sum \frac{\epsilon}{2} \frac{\partial^2}{\partial x_i \partial x_j} (a_{ij} f) - \sum \frac{\partial}{\partial x_i} (b_i f) - \psi f , \quad (4-1)$$

with the following assumptions:

$$A1) \quad a_{ij} = \delta_{ij} = \begin{cases} 1 & \text{if } i=j \\ 0 & \text{otherwise} \end{cases}$$

A2) $b_i(x, t) = b_i$ a constant independent of position and time.

A3) The target moves in the plane.

A4) $\psi(x, t; z) = \bar{\psi}$, a constant.

The search equation is then:

$$\frac{\partial f}{\partial t} = \frac{\epsilon}{2} \left[\frac{\partial^2 f}{\partial x_1^2} + \frac{\partial^2 f}{\partial x_2^2} \right] - \left[b_1 \frac{\partial f}{\partial x_1} + b_2 \frac{\partial f}{\partial x_2} \right] - \bar{\psi} f , \quad (4-2)$$

with

$$f(x, 0; z) = \rho_0(x) \quad (4-3)$$

Exercise

Show that if one sets $f(x, t; z) = w(x, t)e^{-\psi t}$, then:

$$\frac{\partial w}{\partial t} = \frac{c}{2} \left[\frac{\partial^2 w}{\partial x_1^2} + \frac{\partial^2 w}{\partial x_2^2} \right] - \left[b_1 \frac{\partial w}{\partial x_1} + b_2 \frac{\partial w}{\partial x_2} \right] \quad (4-4)$$

with $w(x, 0) = \rho_0(x)$.

CANONICAL PROBLEM: TYPE 1 INITIAL DATA

Assume that:

$$\rho_0(x) = \delta(x) \quad (4-5)$$

i.e., $\Pr\{X(0) = 0\} = 1$.

Then,

$$w(x, t) = \frac{1}{2\pi c t} \exp \left\{ - \left[\frac{(x_1 - b_1 t)^2 + (x_2 - b_2 t)^2}{2c t} \right] \right\}. \quad (4-6)$$

The right-hand side of equation 4-6 is the Green's function, $G(x, t)$, or fundamental solution. As $t \rightarrow 0$

$$w(x, t) \rightarrow \delta(x) = \delta(x_1) \delta(x_2). \quad (4-7)$$

Exercise

Show that equation 4-6 satisfies equation 4-4.

CANONICAL PROBLEM: GENERAL INITIAL DATA

Assume that:

$$\rho_0(x) = \bar{\rho}(x_1, x_2) .$$

Then (see references 5, 14, or 15)

$$\begin{aligned} w(x, t) &= G * \rho \\ &= \int G(x - \xi, t) \rho_0(\xi) d\xi . \end{aligned} \quad (4-8)$$

Exercise

Assume that:

$$\rho_0(x) = \frac{1}{2\pi\sigma^2} \exp\left(-\frac{x_1^2+x_2^2}{2\sigma^2}\right) .$$

Find $w(x, t)$.

Hint: Gaussian functions are closed under convolution
(complete the square).

CANONICAL PROBLEM: TYPE 3 INITIAL DATA

Assume that:

$$\rho_0(x) = \begin{cases} \bar{\rho}(x) & \text{in square of side } 2l \\ 0 & \text{outside} \end{cases} \quad (4-9)$$

Then:

$$w(x, t) = \int_{-l}^l \int_{-l}^l G(x - \xi, t) \bar{\rho}(\xi) d\xi \quad (4-10)$$

Remark

The one-dimensional analogue of equation 4-9 is:

$$\rho_0(x) = \begin{cases} \bar{\rho}(x) & -l \leq x \leq l \\ 0 & \text{otherwise,} \end{cases} \quad (4-11)$$

and of equation 4-11, is

$$w(x, t) = \frac{1}{\sqrt{2\pi\epsilon t}} \int_{-l}^l \bar{\rho}(\xi) \exp\left[-\frac{(x-bt-\xi)^2}{2\epsilon t}\right] d\xi. \quad (4-12)$$

(Hard) Exercise (see references 17 and 18)

Use integration by parts to develop an asymptotic expansion of equation 4-12 valid for small ϵ and $|x-bt| < l$.

Remark

All of the above results also hold when $b_i(x, t) = b_i(t)$ only, independent of x . In this case, the Green's function $G(x, t)$ is:

$$G(x, t) = \frac{1}{2\pi\epsilon t} \exp \left\{ -\frac{1}{2\epsilon t} \left[\left(x_1 - \int_0^t b_1(s) ds \right)^2 + \left(x_2 - \int_0^t b_2(s) ds \right)^2 \right] \right\}. \quad (4-13)$$

CANONICAL PROBLEM: NONCONSTANT ψ

Allow ψ to vary, but keep all the other assumptions. The search equation can be rewritten as:

$$\frac{\partial f}{\partial t} - \frac{\epsilon}{2} \left[\frac{\partial^2 f}{\partial x_1^2} + \frac{\partial^2 f}{\partial x_2^2} \right] + \left[b_1 \frac{\partial f}{\partial x_1} + b_2 \frac{\partial f}{\partial x_2} \right] = -\psi f \equiv u(x, t). \quad (4-14)$$

The solution of equation 4-14 is formally

$$f(x, t; z) = G * u. \quad (4-15)$$

But, $u = -\psi f$, so that:

$$\begin{aligned} f(x, t; z) &= G * (-\psi f) \\ &= -G * (\psi * (G * u)) \\ &\text{etc.} \end{aligned} \quad (4-16)$$

The iterative procedure of equation 4-16 and a suitable fixed point theorem can probably be used to prove the existence of solutions to equation 4-14.

CANONICAL PROBLEM: ORNSTEIN-UHLENBECK PROCESS

Assume that the target moves in one dimension with:

$$\left. \begin{array}{l} b(x,t) = -\beta x \\ a(x) = 1 \end{array} \right\} \quad (4-17)$$

and $\psi = \bar{\psi}$, a constant. Then, if $f(x,t;z) = w(x,t)e^{-\bar{\psi}t}$,
 $w(x,t)$ satisfies:

$$\frac{\partial w}{\partial t} = \frac{c}{2} \frac{\partial^2 w}{\partial x^2} + \beta \frac{\partial}{\partial x} (wx) . \quad (4-18)$$

Assume that $w(x,0) = \delta(x-x_0)$. Feller (reference 5) has shown that:

$$w(x,t) = \left[\frac{\beta}{\pi c (1-e^{-2\beta t})} \right]^{\frac{1}{2}} \exp \left[\frac{-\beta (x-x_0 e^{-2\beta t})^2}{c (1-e^{-2\beta t})} \right] \quad (4-19)$$

Remark

All of these problems have solutions of the form:

$$f \sim w(x,t)e^{-\bar{\psi}t} = [\text{term due to target motion}] \times [\text{term due to search}] .$$

APPROXIMATE SOLUTION OF THE STOCHASTIC
SEARCH EQUATION IN THE GENERAL CASE

General references: 1, 6, 10, 13, 14, 16, 17.

Reconsider the stochastic search equation:

$$\frac{\partial f}{\partial t} = \frac{\epsilon}{2} \sum \frac{\partial^2}{\partial x_i \partial x_j} (a_{ij} f) - \sum \frac{\partial}{\partial x_i} (b_i f) - \psi f , \quad (5-1)$$

with initial data:

$$f(x, 0; z) = p_0(x) . \quad (5-2)$$

For simplicity, assume that the initial data is exponential:

$$p_0(x) = e^{-\phi(x)/\epsilon} \sum_{k=0} \epsilon^k h_k(x) \quad (5-3)$$

Exercise

Cohen and Lewis (reference 13) have shown that the following "trick" can be used to obtain data in the form of equation 5-3.

Show that formally one can set:

$$\begin{aligned} \phi(x) &= -\epsilon \ln p_0(x) \\ h_0(x) &= 1 \\ h_k(x) &\equiv 0 \quad k \geq 1 . \end{aligned} \quad (5-4)$$

For

$$\rho_0(x) = \frac{1}{2\pi} e^{-\frac{(x_1^2+x_2^2)}{2\sigma^2}} \quad (5-5)$$

find $\phi(x)$ and the $g_k(x)$.

Hint: Redefine σ , in terms of ϵ .

Based on the solutions of the canonical problems, one seeks a solution of the SSE in the form (note ϕ does not depend on $z(t)$):

$$f(x,t;z) = e^{-\phi(x,t)/\epsilon} \sum_{k=0} \epsilon^k g_k(x,t;z) . \quad (5-6)$$

In this equation, $\phi(x,t)$ and $g_k(x,t;z)$ $k = 0, 1, 2, \dots$ are to be determined. Equation 5-6 is called a "ray ansatz."

Now apply the following procedure:

1. Evaluate $\partial f / \partial t$, $\partial f / \partial x_i$ and $\partial^2 f / \partial x_i \partial x_j$ if $f(x,t;z)$ is given by equation 5-6.
2. Substitute into the search equation equation 5-1.
3. Collect terms according to powers of ϵ . One obtains:

$$e^{-\phi(x,t)/\epsilon} \left[\frac{1}{\epsilon} \{ \} + \epsilon^0 \{ \} + \epsilon^1 \{ \} + \dots \right] = 0. \quad (5-7)$$

4. Set each coefficient of ϵ^k equal to zero. This gives an equation for $\phi(x,t)$ if $k=-1$ and for $g_k(x,t;z)$ if $k=0, 1, 2, \dots$

Exercise

Show that the equation that $\phi(x,t)$ satisfies is:

$$\frac{\partial \phi}{\partial t} + \sum_i b_i \frac{\partial \phi}{\partial x_i} + \frac{1}{2} \sum_{i,j} a_{ij} \frac{\partial \phi}{\partial x_i} \frac{\partial \phi}{\partial x_j} = 0. \quad (5-8)$$

Equation 5-8 is the "eikonal" or Hamilton-Jacobi equation. It is first order, but non-linear. Let:

$$H(x,p,t) = \sum_i b_i p_i + \frac{1}{2} \sum a_{ij} p_i p_j, \quad (5-9)$$

be a "Hamiltonian" in which $p = (p_1, p_2, p_3)$ is a new independent variable. Then one writes:

$$\frac{\partial \phi}{\partial t} + H\left(x, \frac{\partial \phi}{\partial x}, t\right) = 0. \quad (5-10)$$

Equation (5-8) can also be solved by the method of characteristics.
(references 14, 15, and 16)

Theorem (reference 14,15)

Let $x(t)$ and $p(t)$ be the solutions of:

$$\left. \begin{aligned} \frac{dx_i}{dt} &= \frac{\partial H}{\partial p_i}, & x_i(0) &= x_{i0} \\ \frac{dp_i}{dt} &= -\frac{\partial H}{\partial x_i}, & p_i(0) &= p_{i0} \end{aligned} \right\} \quad (5-11)$$

These are called the ray equations. On the solution curves of equation 5-11:

$$\frac{d\phi}{dt} = -H(x, p_i, t) + \sum p_i \frac{dx_i}{dt} + \frac{1}{2} \sum a_{ij} p_i p_j \quad (5-12)$$

$$\phi(0) = \phi_0$$

Exercise

- 1) Assume that:

$$(a_{ij}) = \begin{pmatrix} 1 & 0 \\ 0 & 1 \end{pmatrix}$$

and $b_i(x, t) = b_i$, a constant. Write down and solve the ray equations.

- 2) Assume that

$$(a_{ij}) = \begin{pmatrix} 1 & 0 \\ 0 & 1 \end{pmatrix}$$

and $b_i(x, t) = b_i(t)$ only. Write down the ray equations. When can they be solved?

- 3) What are the ray equations for general (a_{ij}) and $b_i(x, t)$?

Theorem (references 14 and 15)

If the solution curves of equation 5-11 don't intersect, then $p_i = \partial\phi/\partial x_i$ and $\phi(x, t)$, the solution of equation 5-12, is also a solution of equation 5-8.

Initial conditions are still needed. In light of equations 5-3 and 5-6, one has:

$$e^{-\phi(x)/\epsilon} \sum_k \epsilon^k h_k(x) = e^{-\phi(x,0)/\epsilon} \sum_k g_k(x,0,z) \epsilon^k . \quad (5-13)$$

Hence, one sets:

$$\begin{aligned} x(0) &= x_0 \\ p_i(0) &= \left. \frac{\partial \phi}{\partial x_i} \right|_{x_0} \\ \phi(x_0, 0) &= \phi(x_0) \end{aligned} \quad (5-14)$$

Thus, the following algorithm can be used.

ALGORITHM FOR SOLUTION OF THE HAMILTON-JACOBI EQUATION

1. Pick x_0 (arbitrary).

2. Solve:

$$\begin{aligned} \frac{dx_i}{dt} &= \frac{\partial H}{\partial p_i} & x_i(0) &= x_{i0} \\ \frac{dp_i}{dt} &= -\frac{\partial H}{\partial x_i} & p_i(0) &= \left. \frac{\partial \phi}{\partial x_i} \right|_{x_0} \end{aligned} \quad (5-15)$$

3. Set $\phi(x,0) = \phi_0 = \phi(x_0)$ and solve:

$$\frac{d\phi}{dt} = \frac{1}{2} \sum a_{ij} p_i p_j . \quad (5-16)$$

4. Cycle through x_0 .

ALTERNATIVE ALGORITHM FOR THE SOLUTION OF THE HAMILTON-JACOBI EQUATION

The following alternative algorithm can also be used.

1. Choose a point x at time t .
2. Find the point x_0 such that the solution of

$$\frac{dx_i}{dt} = \frac{\partial H}{\partial p_i} \quad x_i(0) = x_{i0}$$

$$\frac{dp_i}{dt} = - \frac{\partial H}{\partial x_i} \quad p_i(0) = \left. \frac{\partial \phi}{\partial x_i} \right|_{x_0}$$

passes through the point x at time t .

3. Set $\phi_0 = \phi(x_0)$ and solve

$$\frac{d\phi}{dt} = \frac{1}{2} \sum a_{ij} p_i p_j$$

This generates $\phi(x, t)$.

4. Cycle through x at time t .

Variational Interpretation (references 6, 10, 14, 16)

Define a Lagrangian $L(x, dx/dt)$ by:

$$L\left(x, \frac{dx}{dt}, t\right) + H(x, p, t) = \sum_i \frac{dx_i}{dt} p_i . \quad (5-17)$$

Exercise

For $H(x, p)$ given by equation 5-9, show that:

$$L\left(x, \frac{dx}{dt}, t\right) = \frac{1}{2} \sum_{i,j} \left(\frac{dx_i}{dt} - b_i \right) (a_{ij})^{-1} \left(\frac{dx_j}{dt} - b_j \right) . \quad (5-18)$$

In this equation, $(a_{ij})^{-1}$ is the inverse of (a_{ij}) :

$$(a_{ij})(a_{ij})^{-1} = \begin{pmatrix} 1 & 0 \\ 0 & 1 \end{pmatrix} .$$

Let

$$C\{t, x\} = \{ \text{set of all paths } u(s) \text{ with } u(0)=x_0 \text{ and } u(t)=x \}$$

(see figure 3). Then, according to Hamilton's principle:

$$\phi(x, t) = \min_{C\{t, x\}} \int_0^t L\left(u, \frac{du}{ds}, s\right) ds . \quad (5-19)$$

This formulation is often very useful for numerical evaluation of $\phi(x, t)$.




FIG. 3: ILLUSTRATING HAMILTON'S PRINCIPLE

Remarks

1. If there is a deterministic trajectory connecting $(x_0, 0)$ and (x, t) , then $L(x, \frac{dx}{dt}, t)$ vanishes on that trajectory.
2. If $\rho_0(x) = \delta(x - x_0)$ then the p_i 's can be treated as parameters (see reference 1 or 6).
3. If $\rho_0(x)$ is type 3, i.e.:

$$\rho_0(x) = \begin{cases} \bar{\rho}(x) & x \in D_T' \cap D_T \\ 0 & x \notin D_T' \end{cases}$$

there may be difficulties with the ansatz (equation 5-6) (i.e., rays may intersect for x_0 near the boundary of D_T'). More work (boundary layer solutions or uniform methods) can eradicate the difficulties.

Now consider $g_0(x, t; z)$.

Exercise

Show that $g_0(x, t; z)$ satisfies the following equation:

$$\frac{\partial g_0}{\partial t} + \sum_i \left(\sum_j a_{ij} \frac{\partial \phi}{\partial x_j} + b_i \right) \frac{\partial g_0}{\partial x_i}$$

(5-20)

$$= - \left[\sum_{i,j} \left(\frac{\partial a_{ij}}{\partial x_i} \frac{\partial \phi}{\partial x_j} + \frac{1}{2} a_{ij} \frac{\partial^2 \phi}{\partial x_i \partial x_j} \right) + \sum_i \frac{\partial b_i}{\partial x_i} + \psi(x, t; z) \right] g_0 .$$

This is a first-order linear differential equation for $g_0(x, t; z)$; it is called a transport equation. Note that if $(a_{ij}) = (0)$, then equation 5-20 becomes the deterministic search equation.

To solve equation 5-20, recall that:

$$\frac{dx_i}{dt} = \frac{\partial H}{\partial p_i} = b_i + \sum_j a_{ij} \frac{\partial \phi}{\partial x_j} . \quad (5-21)$$

Then equation 5-20 becomes:

$$\begin{aligned} \frac{\partial g_0}{\partial t} + \sum_i \frac{dx_i}{dt} \frac{\partial g_0}{\partial x_i} &= - \left[\sum_{i,j} \left(\frac{\partial a_{ij}}{\partial x_i} p_j + \frac{1}{2} a_{ij} \frac{\partial^2 \phi}{\partial x_i \partial x_j} \right) \right. \\ &\quad \left. + \sum_i \frac{\partial b_i}{\partial x_i} + \psi(x, t; z) \right] g_0 . \end{aligned} \quad (5-22)$$

Thus, on the rays:

$$\frac{dg_0}{dt} = - \{ r(t) + \psi(x(t), t, z) \} g_0 . \quad (5-23)$$

In this equation:

$$r(t) = \sum_{i,j} \left(\frac{\partial a_{ij}}{\partial x_j} p_j + \frac{1}{2} a_{ij} \frac{\partial^2 \phi}{\partial x_i \partial x_j} \right) + \sum_i \frac{\partial b_i}{\partial x_i} . \quad (5-24)$$

An alternative formulation is obtained by following reference 6.

Let $J(t)$ satisfy:

$$\frac{dJ}{dt} = J \sum_i \frac{\partial}{\partial x_i} \left(\frac{dx_i}{dt} \right) . \quad (5-25)$$

Then equation 5-24 becomes:

$$\Gamma(t) = \frac{1}{2J} \frac{dJ}{dt} + \frac{1}{2} \sum_{i,j} \frac{\partial a_{ij}}{\partial x_j} \frac{\partial \phi}{\partial x_i} + \sum_i \frac{\partial b_i}{\partial x_i} , \quad (5-26)$$

so that $\partial^2 \phi / \partial x_i \partial x_j$ does not have to be computed.

Remark

Ludwig (reference 6) interprets J as the Jacobian of the transformation from physical to "ray space."

Initial Condition

From equation 5-3 and 5-6, it is clear the the appropriate initial data is:

$$g_0(x_0, 0; z) = h_0(x_0) . \quad (5-27)$$

The other functions $g_k(x, t; z)$ are determined in a similar fashion.

SUMMARY OF COMPUTATIONAL ALGORITHMS

In this section, three algorithms are given for the solution of the three search equations corresponding to different types of target motion.

DETERMINISTIC TARGET MOTION

Equation:

$$\frac{\partial f}{\partial t} = - \sum_i \frac{\partial}{\partial x_i} (b_i f) - \psi f$$

$$f(x, 0; z) = p_0(x)$$

Algorithm

1. Pick x_0

2. Solve

$$\frac{dx_i}{dt} = b_i(x, t) \quad x_i(0) = x_{i0}$$

3. Solve

$$\frac{df}{dt} = - \left[\sum_i \frac{\partial b_i(x(t), t)}{\partial x_i} + \psi(x(t), t, z) \right] f$$

with $f(0) = p_0(x_0)$

4. Cycle to 1 or 2.

CONDITIONALLY DETERMINISTIC TARGET MOTION

Equation:

$$\frac{\partial f}{\partial t} = - \sum_i \frac{\partial}{\partial x_i} \left(b_i^2 f \right) - \psi f \quad (1)$$

$$f(x, 0; z) = p_0(x)$$

$b_\alpha(x, t)$ drawn from a distribution $B(b_\alpha)$.

Algorithm

1. Fix α and let $f(x, t; z|\alpha)$ denote the solution of (1) with α fixed.
2. Obtain $f(x, t; z|\alpha)$ by using steps 1 through 4 of the previous algorithm.
3. Calculate:

$$f(x, t; z) = E_\alpha \{ f(x, t; z|\alpha) \} = \int_{B(b_\alpha)} f(x, t; z|\alpha) dP_\alpha.$$

STOCHASTIC TARGET MOTION

Equation:

$$\frac{\partial f}{\partial t} = \frac{\epsilon}{2} \sum_{i,j} \frac{\partial^2}{\partial x_i \partial x_j} (a_{ij} f) - \sum_i \frac{\partial}{\partial x_i} (b_i f) - \psi f$$

$$f(x, 0; z) = e^{-\Phi(x)/\epsilon} \sum_{k=0} \epsilon^k h_k(x).$$

Algorithm for $f(x,t,z) \sim e^{-\phi(x,t)} / c_{g_0}(x,t,z)$

1. Pick x at t ; find x_0 .

2. Find $p_{i0} = \left. \frac{\partial \phi}{\partial x_i} \right|_{x=x_0}$

3. Solve the ray equations:

$$\frac{dx_i}{dt} = \frac{\partial H}{\partial p_i} \quad x_i(0) = x_{i0}$$

$$\frac{\partial p_i}{dt} = - \frac{\partial H}{\partial x_i} \quad p_i(0) = p_{i0}$$

where

$$H(x, p, t) = \sum b_i p_i + \frac{1}{2} \sum a_{ij} p_i p_j .$$

4. Solve for ϕ :

$$\frac{d\phi}{dt} = \frac{1}{2} \sum_{i,j} a_{ij} p_i p_j$$

$$\phi(0) = \phi(x_0)$$

5. Evaluate the Jacobian $J(t)$ or calculate $\partial^2 \phi / \partial x_i \partial x_j$

6. Solve for g_0 :

$$\frac{dg_0}{dt} = - [r(t) + \psi(x, t, z)] g_0$$

$$g_0(x_0, 0, z) = h_0(x_0)$$

7. Cycle to 1.

OPTIMAL SEARCH FOR A MOVING OBJECT

General References: 2, 3, 17.

In this section, the problem of optimal search is formulated and a method of solution is sketched.

Let $C(z)$ be the set of allowable search paths. For example:

$$C(z) = \{z(t): z(0)=z_0, z(T_f)=z_f, \left| \frac{dz}{dt} \right| \leq v, \left| \frac{d^2z}{dt^2} \right| \leq a\}, \quad (7-1)$$

is the set of search paths with constrained endpoints and constraints on velocity and acceleration.

Three possible objective functionals are the following.
(Other functionals could be used.)

FUNCTIONAL #1: INSTANTANEOUS PROBABILITY OF DETECTION

Let:

$$J_1(z,t) = E_x\{\psi(x,t,z)\} \quad (7-2)$$

$$= \int \psi(x,t,z) \rho(x,t|z) dx . \quad (7-3)$$

Note that $J_1(z,t)$ is the (averaged) instantaneous probability of detection. A search plan that maximizes $J_1(z,t)$ is called myopic.

From the definition of $\rho(x,t|z)$, one obtains that:

$$J_1(z,t) = \frac{\int \psi(x,t,z) f(x,t;z) dx}{\int f(x,t;z) dx} . \quad (7-4)$$

When $J_1(z,t)$ is extremized, one has the following interpretation.

Consider an interval $(0,T)$. Divide it into k subintervals

(t_i, t_{i+1}) with $t_0 = 0$ and $t_k = T$.

Next, maximize $J_1(z,t_i)$ for $i = 1, 2, \dots, k-1$. The optimal path is the piecewise linear set $[z_0, z_1], [z_1, z_2]$, etc., where

$$z_i = z(t_i).$$

As $k \rightarrow \infty$, a continuous curve $z^*(t)$ is obtained.

FUNCTIONAL #2: PROBABILITY OF DETECTION BY TIME T_S

Let

$$J_2(z, T_S) = 1 - \int f(x, T_S; z) dx . \quad (7-5)$$

This functional, $J_2(z, T_S)$, is the probability of detection by time T_S . One seeks the path $z^*(t)$, $0 \leq t \leq T_S$ that maximizes $J_2(z, T_S)$. Also, note that $f(x, t; z)$ satisfies:

$$\frac{\partial f}{\partial t} = \frac{1}{2} \sum_{i,j} \frac{\partial^2}{\partial x_i \partial x_j} (a_{ij} f) - \sum_i \frac{\partial}{\partial x_i} (b_i f) - \psi f . \quad (7-6)$$

In terms of control theory, this is an optimal control problem for a distributed parameter system with control in the coefficients. There is essentially no theory for this sort of problem.

FUNCTIONAL # 3: MEAN TIME TO DETECTION

The probability density for the time of detection can be determined as follows. Let T_F be the total time available for search. Let $h(t)dt = \Pr\{\text{detection occurs in } (t, t+dt)\}$. Then:

$$h(t) = \begin{cases} -\frac{1}{T_F} \int f(x, t; z) dx & t \leq T_F \\ 0 & t > T_F \end{cases} \quad (7-7)$$

is the desired probability density.

The mean time to detection is then:

$$J_3(z) = \int_0^{T_F} th(t)dt \quad (7-8)$$

$$= \int_0^{T_F} t \left[-\frac{1}{T_F} \int f(x, t; z) dx \right] dt \quad (7-9)$$

SOLUTION OF AN OPTIMAL SEARCH PROBLEM

Consider now the problem of finding $z^*(\tau) \in C(z)$ that maximizes $J_2(z, T_S)$. In order to make any progress, the functional will be converted to a function as follows. The functional $J_2(z, t)$ can be rewritten as:

$$\begin{aligned} J_2(z; t) &= 1 - \int e^{-\phi(x, t)/\epsilon} \sum \epsilon^k g_k(x, t; z) dx \\ &= 1 - \sum \epsilon^k \int e^{-\phi(x, t)/\epsilon} g_k(x, t; z) dx \end{aligned} \quad (7-10)$$

$$= 1 - \sum_{k=0} \epsilon^k I_k(z, t) . \quad (7-11)$$

The integrals $I_k(z, t)$ are defined by:

$$I_k(z, t) = \int e^{-\psi(x, t)/\epsilon} g_k(x, t; z) dx . \quad (7-12)$$

These integrals can be analyzed by Laplace's method (reference 17).

Let $x^*(t)$ denote the minimum over D_T of $\psi(x, t)$. For simplicity, assume that there is only one minimum. From reference 17, page 338, one has that:

$$I_k(z, t) \sim e^{-\psi(x^*(t), t)/\epsilon} (2\pi\epsilon)^{n/2} \sum_{j=0}^{\infty} \frac{\Delta_\epsilon^j G_{k0}}{j! 2^j} \Big|_{x=x^*} . \quad (7-13)$$

In this equation, $n = 2$ or 3 is the dimension of the space in which the target moves. The variables ξ_i are now coordinates (reference 17, page 334),

$$\Delta_\epsilon^j = \left(\sum_{i=1} \frac{\partial^2 \psi}{\partial \xi_i^2} \right)^j \quad (7-14)$$

and

$$G_{k0} = \frac{g_k(x^*(t), t; z)}{\left[\det \left(\frac{\partial^2 \psi}{\partial \xi_i \partial \xi_j} \right) \Big|_{x^*(t)} \right]^{\frac{1}{2}}} \quad (7-15)$$

It is often sufficient to use only the first one or two terms of the full asymptotic expansion.

Thus, $J_2(z, T_S)$ has now been converted to a function, rather than functional:

$$J_2(z, T_S) = 1 - (2\pi\epsilon)^{n/2} e^{-\phi(x^*(T_S), T_S)/\epsilon} \quad (7-16)$$

$$\sum_{k=0}^{\infty} \sum_{j=0}^{\infty} \epsilon^k \left[\sum_{j=0}^{\infty} \frac{\Delta_j^j G_{k0}}{j!} \left(\frac{\epsilon}{2}\right)^j \right] .$$

One now determines the function $z^*(t)$ that maximizes equation 7-14 on $[0, T_S]$. Here, $z(t)$ occurs in the G_{k0} -terms, which depend on $g_k(x, t; z)$ by equation 7-15, and the dependence of g_k on $z(t)$ is given by equations such as 5-23.

Remarks

1. It is still, in most cases, necessary to use a computer to obtain $z^*(t)$. However, by going from functional to function, the amount of work required has been vastly reduced.
2. It may be possible to solve some of these problems by "dynamic programming."

REFERENCES

1. Center for Naval Analyses, Memorandum (CNA)78-1365, "New Methods in Search for a Moving Target," Marc Mangel, Unclassified, 11 Sep 1978.
2. Center for Naval Analyses, OEG Report 56, "Search and Screening," B.O. Koopman, Unclassified , 1946.
3. Stone, L., "Theory of Optimal Search," Academic Press, New York, 1975.
4. Corwin, T., "An Expression for the Location Density of a Moving Target in a Multiple Sensor Environment," D.H. Wagner and Associates, Memorandum 1975.
5. Feller, W. , "An Introduction to Probability Theory and Its Applications," Wiley & Sons, New York, 1965.
6. Ludwig, D., "Persistence of Dynamical Systems Under Random Perturbations," SIAM Review, 1975, 17:605-635.
7. Arnold, L., "Stochastic Differential Equations," Wiley & Sons, New York, 1974.
8. Lighthill, M.J., "Generalized Functions," Cambridge University Press, Cambridge, 1975.
9. U.S.N.A., "Naval Operations Analysis" 1975.
10. Ludwig, D., "Stochastic Population Theories," Springer Verlag, New York, 1974.
11. Hellman, O., "On the Effect of a Search Upon the Probability Density of a Target Whose Motion is a Diffusion Process," Annals of Mathematical Statistics, 1970, 41:1717-1724.
12. Ciervo, A., "Search for Moving Targets," Pacific Sierra Research Corporation, Report 619B, 1976.
13. Cohen, J.K. and R.M. Lewis, "A Ray Method for the Asymptotic Solution of the Diffusion Equation," Journal Institute of Mathematical Applications, 1967, 3: 226-290.
14. Courant, R. and D. Hilbert, "Methods of Mathematical Physics, Volume II," Wiley & Sons, New York, 1962.

15. Carrier, G.F. and C. Pearson, "Partial Differential Equations,"
16. Keller, J.B., "Rays, Waves, and Asymptotics," Bulletin of American Mathematical Society, 1978, 84: 727-750.
17. Bleistein, N. and R.A. Handelsman, "Asymptotic Expansion of Integrals," Irvington, New York, 1978.
18. Olver, F.J.W., "Asymptotics and Special Functions," Academic, New York, 1974.

APPENDIX A
SOLUTIONS OF EXERCISES

In this appendix, written by James Thomas, Jr., the solutions of most of the exercises are given.

EXERCISE:

$$\delta(s) = \lim_{n \rightarrow \infty} \frac{\sqrt{n}}{\sqrt{2\pi}} e^{-s^2 n/2}$$

$$= \lim_{n \rightarrow \infty} \xi_n(s)$$

$$\text{with } \xi_n(s) = \frac{\sqrt{n}}{\sqrt{2\pi}} e^{-s^2 n/2} .$$


Sketch a few $\xi_n(s)$.

Sketch the corresponding $\xi'_n(s)$, where the prime indicates a derivative with respect to s .

SOLUTION:


Refer to the following computer graphics.

$$f_n = \sqrt{\frac{n}{2\pi}} e^{-ns^2/2}$$



$$\xi_n' = -\frac{n^{3/2}}{\sqrt{2\pi}} s e^{-ns^2/2}$$

$$\xi_n' = -ns\xi_n$$



EXERCISE:

Show that for deterministic target motion:

$$\text{var}\{\Delta X | X(t)=x\} = 0.$$

SOLUTION:

For deterministic target motion, the transition function,
 $q(\xi, t, \Delta t, x)$ is given by

$$q(\xi, t, \Delta t, x) = \delta(\xi - b(x, t)\Delta t).$$

Substitution of this in the expression,

$$E\{(\Delta X)^2\} = \int \xi^2 q d\xi ,$$

yields

$$E\{(\Delta X)^2\} = \int \xi^2 \delta(\xi - b(x, t)\Delta t) d\xi = [b(x, t)]^2 (\Delta t)^2.$$

Consequently,

$$\text{var}\{\Delta X\} = E\{(\Delta X)^2\} - [E\{\Delta X\}]^2 = 0 .$$

For stochastic target motion, given $x(t) = x$, ΔX is distributed with mean

$$b(x, t)\Delta t + o(\Delta t)$$

and covariance

$$\epsilon a_{ij}(x, t)\Delta t + o(\Delta t),$$

where

$$\frac{o(\Delta t)}{\Delta t} \rightarrow 0 \quad \text{as } \Delta t \rightarrow 0.$$

Alternatively:

$$E\{\Delta X_i | X(t)=x\} = b_i(x, t)\Delta t + o(\Delta t)$$

and

$$E\{\Delta X_i \Delta X_j | X(t)=x\} = \epsilon a_{ij}\Delta t + o(\Delta t).$$

EXERCISE:

If ΔX is normally distributed, what happens as $\epsilon \rightarrow 0$ with a_{ij} bounded? Write the alternative conditions for stochastic target motion in terms of q .

SOLUTION: Let $dX = X(t + dt) - X(t)$.

In several dimensions,

$$q(\xi_1, \xi_2, \dots, \xi_j, t, dt, x) d\xi_1 d\xi_2 \dots d\xi_j$$

$$= \text{Prob}\{\xi_1 \leq X_1(t+dt) - X_1(t) \leq \xi_1 + d\xi_1, \xi_2 \leq X_2(t+dt) - X_2(t) \leq \xi_2 + d\xi_2,$$

$$\dots, \xi_j \leq X_j(t+dt) - X_j(t) \leq \xi_j + d\xi_j \mid X(t) = x\}.$$

If two new functions are defined such that

$$q_{ij}(\xi_i, \xi_j, t, dt, x) = \int \dots \int d\xi_1 d\xi_2 \dots d\xi_{i-1} q(\xi_1, \xi_2, \dots, \xi_i, \xi_j, t, dt, x)$$

and

$$q_i(\xi_i, t, dt, x) = \int \dots \int d\xi_1 d\xi_2 \dots d\xi_{i-1} d\xi_j q(\xi_1, \xi_2, \dots, \xi_i, \xi_j, t, dt, x),$$

then the alternative conditions for stochastic target motion can be written

$$\begin{aligned} E(dx_i \mid X(t) = x) &= b_i(x, t) dt + o(dt) \\ &= \int d\xi_i q_i(\xi_i, t, dt, x) \xi_i \end{aligned}$$

and

$$\begin{aligned} E(dx_i dx_j \mid X(t) = x) &= \epsilon a_{ij} dt + o(dt) \\ &= \iint d\xi_i d\xi_j q_{ij}(\xi_i, \xi_j, t, dt, x) \xi_i \xi_j. \end{aligned}$$

Now, given that dX is normally distributed with mean,

$$b(x, t) dt + o(dt),$$

and covariance,

$$\epsilon a_{ij}(x,t)dt + o(dt),$$
$$q_{ij}(\xi_i, \xi_j, t, dt, x) = \frac{1}{\sqrt{2\pi[\epsilon a_{ij}dt + o(dt)]}}$$
$$\cdot \exp \left\{ -\frac{[\xi_i - b_i dt - o(dt)][\xi_j - b_j dt - o(dt)]}{2[\epsilon a_{ij}dt + o(dt)]} \right\}$$

As $\epsilon \rightarrow 0$ with a_{ij} bounded, $q_{ij}(\xi_i, \xi_j, t, dt, x)$ becomes a delta function analogous to the $\delta_n(s)$ of the first exercise. Consequently, the target motion becomes deterministic in x , i.e.,

$$dx_i = b_i(x,t)dt \quad \text{with probability 1}.$$

EXERCISE:

For one-dimensional target motion with $dX \sim N(b(x,t)dt + o(dt); \epsilon a(x,t)dt + o(dt))$, show that

$$\int \xi^n q(\xi, t, dt, x) d\xi = o(dt)$$

for $n \geq 3$, where

$$\frac{o(dt)}{dt} \rightarrow 0 \quad \text{as } dt \rightarrow 0$$

SOLUTION:

$$\text{For } dX \sim N(b(x,t)dt + o(dt); \epsilon a(x,t)dt + o(dt)),$$

$$q(\xi, t, dt, x) = \frac{1}{\sqrt{2\pi} [\epsilon a(x,t)dt + o(dt)]} e^{-\frac{[\xi - b(x,t)dt - o(dt)]^2}{2[\epsilon a(x,t)dt + o(dt)]}}$$

For convenience, let

$$q(\xi, t, dt, x) = \frac{1}{\sqrt{2\pi} \cdot A} e^{-\frac{[\xi - B]^2}{2A}}$$

with

$$A = \epsilon a(x,t)dt + o(dt)$$

and

$$B = b(x,t)dt + o(dt).$$

Now

$$\int \xi^n q(\xi, t, dt, x) d\xi = \frac{1}{\sqrt{2\pi} A} \int \xi^n e^{-\frac{[\xi - B]^2}{2A}} d\xi.$$

Making the change of variable ,

$$y = \xi - B ,$$

$$\begin{aligned} \int \xi^n q(\xi, t, dt, x) d\xi &= \frac{1}{\sqrt{2\pi A}} \int [y + B]^n e^{-\frac{y^2}{2A}} dy \\ &= \frac{1}{\sqrt{2\pi A}} \int [y^n + Bny^{n-1} + B^2 \frac{n(n-1)}{2} y^{n-2} + \dots] e^{-\frac{y^2}{2A}} dy , \end{aligned}$$

for which only the even powers of y are non-vanishing.

For n even:

$$\begin{aligned} \int \xi^n q(\xi, t, dt, x) d\xi &= \frac{1}{\sqrt{2\pi A}} \int [y^n + B^2 \frac{n(n-1)}{2} y^{n-2} + \dots] e^{-\frac{y^2}{2A}} dy \\ &= \frac{1}{\sqrt{\pi}} \Gamma\left(\frac{n+1}{2}\right) (2A)^{n/2} \left[1 + \frac{nB^2}{(2A)} + \dots \right] \\ &= \frac{2^{n/2}}{\sqrt{\pi}} \Gamma\left(\frac{n+1}{2}\right) (ea)^{n/2} (dt)^{n/2} + \text{terms of higher} \\ &\quad \text{order in } (dt). \end{aligned}$$

Thus, $\int \xi^n q(\xi, t, dt, x) d\xi = o(dt)$ for n even ≥ 4 .

For n odd:

$$\begin{aligned} \int \xi^n q(\xi, t, dt, x) d\xi &= \frac{1}{\sqrt{2\pi A}} \int [nBy^{n-1} + B^3 \frac{n(n-1)(n-2)}{6} y^{n-3} + \dots] e^{-\frac{y^2}{2A}} dy \\ &= \frac{2B}{\sqrt{\pi}} \Gamma\left(\frac{n+2}{2}\right) (2A)^{\frac{(n-1)}{2}} \left[1 + \frac{(n-1)B^2}{3(2A)} + \dots \right] \\ &= \frac{2^{(n+1)/2}}{\sqrt{\pi}} \Gamma\left(\frac{n+2}{2}\right) b \cdot (ea)^{\frac{(n-1)}{2}} (dt)^{\frac{(n+1)}{2}} + \text{terms} \\ &\quad \text{higher order in } (dt). \end{aligned}$$

Thus,

$$\int \xi^n q(\xi, t, dt, x) d\xi = o(dt) \text{ for } n \text{ odd} \geq 3;$$

and

$$\therefore \int \xi^n q(\xi, t, dt, x) d\xi = o(dt) \text{ for } n \geq 3.$$

EXERCISE:

Suppose that there is no search. Let

$$\rho(x, t) dx = \text{Prob}\{X(t) \in dx\} .$$

What equation does ρ satisfy?

SOLUTION:

$$\begin{aligned}
 \rho(\underline{x}, t+dt) &= \int q(\xi, t, dt, \underline{x}-\xi) \rho(\underline{x}-\xi, t) d\xi \\
 &= \int d\xi \left[q\rho - \sum_i \xi_i \frac{\partial}{\partial x_i} (\rho q) + \frac{1}{2} \sum_{i,j} \xi_i \xi_j \frac{\partial^2}{\partial x_i \partial x_j} (\rho q) \right] \\
 \text{where } q\rho &= q(\xi, t, dt, \underline{x}) \rho(\underline{x}, t) \\
 &= \rho(\underline{x}, t) \int d\xi q - \sum_i \frac{\partial}{\partial x_i} \rho \int d\xi \xi_i q + \frac{1}{2} \sum_{i,j} \frac{\partial^2}{\partial x_i \partial x_j} \rho \int d\xi \xi_i \xi_j q .
 \end{aligned}$$

Thus,

$$\begin{aligned}
 \rho(\underline{x}, t+dt) - \rho(\underline{x}, t) &= - \sum_i \frac{\partial}{\partial x_i} \rho [b_i(\underline{x}, t) dt + o(dt)] \\
 &\quad + \frac{1}{2} \sum_{i,j} \frac{\partial^2}{\partial x_i \partial x_j} \rho [\epsilon_{ij}(\underline{x}, t) dt + o(dt)] .
 \end{aligned}$$

Dividing by dt and taking the limit $dt \rightarrow 0$,

$$\frac{\partial \rho}{\partial t} + \sum_i \frac{\partial}{\partial x_i} (b_i \rho) = \frac{1}{2} \sum_{i,j} \frac{\partial^2}{\partial x_i \partial x_j} (\epsilon_{ij} \rho) .$$

EXERCISE:

For deterministic target motion, show that

$$\frac{\partial f}{\partial t} = - \sum_i \frac{\partial}{\partial x_i} (b_i f) = \eta f.$$

SOLUTION:

For deterministic target motion, $\sigma^2 \rightarrow 0$ in the stochastic search equation with a_{ij} bounded.

∴ Search equation becomes

$$\frac{\partial f}{\partial t} = -\eta f - \sum_i \frac{\partial}{\partial x_i} (b_i f) .$$

This result is also obtained by setting

$$q(\xi, t, dt, x) = \delta(\xi - b(x, t)dt) .$$

EXERCISE:

Suppose that $b_i(x, t) = v_i(t)$. Show that

$$f(x, t; z) = p_0 \left(x - \int_0^t v(s) ds \right) \cdot \exp \left[- \int_0^t \psi \left(x - \int_s^t v(s') ds', s; z \right) ds \right].$$

Write down the integral that gives the probability of detection by time t .

SOLUTION:

If $b_i(x, t) = v_i(t)$, $\frac{\partial b_i}{\partial x_i} = 0$, and

$$f(x, t; z) = p_0(x_0) \text{traj}_x \exp \left[- \int_0^t \left(\psi + \sum_i \frac{\partial b_i}{\partial x_i} \right) ds \right]$$

becomes

$$f(x, t; z) = p_0(x_0) \text{traj}_x \exp \left[- \int_0^t \psi \left(x(s), s; z \right) ds \right].$$

Now $x(s) = x_0 + \int_0^s v(s') ds'$,

and $x_0 = x - \int_0^t v(s) ds = x - \int_0^t v(s') ds'$.

Thus,

$$x(s) = x(t) - \int_0^t v(s') ds' + \int_0^s v(s') ds'$$

$$= x(t) - \int_0^t v(s') ds' - \int_s^0 v(s') ds'$$

$$= x(t) - \int_s^t v(s') ds' ,$$

and

$$f(x, t; z) = p_0 \left(x - \int_0^t v(s) ds \right)$$

$$\cdot \exp \left[- \int_0^t \psi \left(x - \int_s^t v(s') ds', s; z \right) ds \right] .$$

$$\text{Prob}\{\text{detection by time } t\} = 1 - \text{Prob}\{\text{no detection}\}$$

$$= 1 - \int f(x, t; z) dx$$

$$= 1 - p_0 \left(x - \int_0^t v(s) ds \right)$$

$$\cdot \exp \left[- \int_0^t \psi \left(x - \int_s^t v(s') ds', s; z \right) ds \right] dx .$$

EXERCISE:

Let $f(x, t) = w(x, t)e^{-\bar{\psi}t}$. Show that

$$\frac{\partial w}{\partial t} = \frac{\epsilon}{2} \sum_{i,j} \delta_{ij} \frac{\partial^2 w}{\partial x_i \partial x_j} - \sum_i b_i \frac{\partial w}{\partial x_i}$$

and $w(x, 0) = \rho_0(x)$.

SOLUTION:

Given $f(x, t) = w(x, t)e^{-\bar{\psi}t}$,

$$\frac{\partial f}{\partial t} = \frac{\partial w}{\partial t} e^{-\bar{\psi}t} - \bar{\psi}w(x, t)e^{-\bar{\psi}t}$$

$$= \left(\frac{\partial w}{\partial t} \right) e^{-\bar{\psi}t} - \bar{\psi}f(x, t).$$

For $a_{ij} = \delta_{ij}$ and b_i constant,

$$\frac{\partial f}{\partial t} = \frac{\epsilon}{2} \sum_{i,j} \frac{\partial^2}{\partial x_i \partial x_j} (a_{ij} f) - \sum_i \frac{\partial}{\partial x_i} (b_i f) - \bar{\psi}f$$

becomes

$$\frac{\partial f}{\partial t} = \frac{\epsilon}{2} \sum_{i,j} \delta_{ij} \frac{\partial^2 f}{\partial x_i \partial x_j} - \sum_i b_i \frac{\partial f}{\partial x_i} - \bar{\psi}f$$

$$= \left(\frac{\partial w}{\partial t} \right) e^{-\bar{\psi}t} - \bar{\psi}f.$$

Thus,

$$\begin{aligned} \left(\frac{\partial w}{\partial t} \right) e^{-\bar{\psi}t} &= \frac{\epsilon}{2} \sum_{i,j} \delta_{ij} \frac{\partial^2 f}{\partial x_i \partial x_j} - \sum_i b_i \frac{\partial f}{\partial x_i} \\ &= \left[\frac{\epsilon}{2} \sum_{i,j} \delta_{ij} \frac{\partial^2 w}{\partial x_i \partial x_j} - \sum_i b_i \frac{\partial w}{\partial x_i} \right] e^{-\bar{\psi}t}, \end{aligned}$$

and $\therefore \frac{\partial w}{\partial t} = \frac{\epsilon}{2} \sum_{i,j} \delta_{ij} \frac{\partial^2 w}{\partial x_i \partial x_j} - \sum_i b_i \frac{\partial w}{\partial x_i}$.

Also, $w_0(x) = f(x, 0) = w(x, 0)$

EXERCISE:

In two dimensions, $w(x,t) = G(x-\xi, t)\rho_0(\xi)d\xi$, with

$$G(x-\xi, t) = \frac{1}{2\pi\epsilon t} \exp\left[-\frac{(x_1-b_1t-\xi_1)^2 + (x_2-b_2t-\xi_2)^2}{2\epsilon t}\right].$$

Assume

$$\rho_0(x) = \frac{1}{2\pi\sigma^2} e^{-\left[\frac{(x_2)^2 + (x_1)^2}{2\sigma^2}\right]}.$$

Find $w(x,t)$. HINT: Gaussians are closed under convolution.

SOLUTION:

$$\begin{aligned} w(x,t) &= \int G(x-\xi, t)\rho_0(\xi)d\xi \\ &= \int d\xi_1 d\xi_2 \frac{1}{2\pi\epsilon t} \exp\left[-\frac{(x_1-b_1t-\xi_1)^2 + (x_2-b_2t-\xi_2)^2}{2\epsilon t}\right] \\ &\quad \cdot \frac{1}{2\pi\sigma^2} \exp\left[-\frac{(\xi_2)^2 + (\xi_1)^2}{2\sigma^2}\right] \\ &= \frac{1}{4\pi^2\sigma^2\epsilon t} \int \exp\left[-\frac{(x_1-b_1t-\xi_1)^2}{2\epsilon t} - \frac{(\xi_1)^2}{2\sigma^2}\right] d\xi_1 \\ &\quad \cdot \int \exp\left[-\frac{(x_2-b_2t-\xi_2)^2}{2\epsilon t} - \frac{(\xi_2)^2}{2\sigma^2}\right] d\xi_2. \end{aligned}$$

Now,

$$(x_1 - b_1 t - \xi_1)^2 = (\xi_1)^2 - 2(x_1 - b_1 t)\xi_1 + (x_1 - b_1 t)^2$$

and

$$\frac{(x_1 - b_1 t - \xi_1)^2}{2\epsilon t} + \frac{(\xi_1)^2}{2\sigma^2} = \frac{2(\sigma^2 + \epsilon t)(\xi_1)^2 - 4\sigma^2(x_1 - b_1 t)\xi_1 + 2\sigma^2(x_1 - b_1 t)^2}{4\sigma^2 \epsilon t}$$

$$= \frac{\left[\sqrt{2(\sigma^2 + \epsilon t)} \xi_1 - \frac{\sqrt{2}\sigma^2(x_1 - b_1 t)}{\sqrt{(\sigma^2 + \epsilon t)}} \right]^2 + \left[\frac{2\sigma^2 \epsilon t (x_1 - b_1 t)^2}{(\sigma^2 + \epsilon t)} \right]}{4\sigma^2 \epsilon t}$$

Consequently,

$$\begin{aligned} & \int \exp \left[-\frac{(x_1 - b_1 t - \xi_1)^2}{2\epsilon t} - \frac{(\xi_1)^2}{2\sigma^2} \right] d\xi_1 \\ &= \int \exp \left\{ - \left[\frac{\sqrt{2}(\sigma^2 + \epsilon t)\xi_1 - \sqrt{2}\sigma^2(x_1 - b_1 t)}{\sqrt{4\sigma^2 \epsilon t} \sqrt{(\sigma^2 + \epsilon t)}} \right]^2 - \left[\frac{(x_1 - b_1 t)^2}{2(\sigma^2 + \epsilon t)} \right] \right\} d\xi_1 \\ &= \exp \left[-\frac{(x_1 - b_1 t)^2}{2(\sigma^2 + \epsilon t)} \right] \int d\xi_1 \exp \left\{ - \left[\frac{\sqrt{2}(\sigma^2 + \epsilon t)\xi_1 - \sqrt{2}\sigma^2(x_1 - b_1 t)}{\sqrt{4\sigma^2 \epsilon t} \sqrt{(\sigma^2 + \epsilon t)}} \right]^2 \right\} \\ &= \exp \left[-\frac{(x_1 - b_1 t)^2}{2(\sigma^2 + \epsilon t)} \right] \frac{\sqrt{2\sigma^2 \epsilon t}}{\sqrt{(\sigma^2 + \epsilon t)}} \int e^{-y^2} dy \end{aligned}$$

$$= \sqrt{\frac{2\pi\sigma^2\epsilon t}{(\sigma^2+\epsilon t)}} \exp\left[-\frac{(x_1-b_1t)^2}{2(\sigma^2+\epsilon t)}\right].$$

Similarly,

$$\int \exp\left[-\frac{(x_2-b_2t-\xi_2)^2}{2\epsilon t} - \frac{(\xi_2)^2}{2\sigma^2}\right] d\xi_2 = \sqrt{\frac{2\pi\sigma^2\epsilon t}{(\sigma^2+\epsilon t)}} \exp\left[-\frac{(x_2-b_2t)^2}{2(\sigma^2+\epsilon t)}\right],$$

and

$$w(x,t) = \frac{1}{4\pi^2\sigma^2\epsilon t} \int \exp\left[-\frac{(x_1-b_1t-\xi_1)^2}{2\epsilon t} - \frac{(\xi_1)^2}{2\sigma^2}\right] d\xi_1$$

$$\cdot \int \exp\left[-\frac{(x_2-b_2t-\xi_2)^2}{2\epsilon t} - \frac{(\xi_2)^2}{2\sigma^2}\right] d\xi_2$$

$$= \frac{1}{2\pi(\sigma^2+\epsilon t)} \exp\left[-\frac{(x_1-b_1t)^2 + (x_2-b_2t)^2}{2(\sigma^2+\epsilon t)}\right].$$

EXERCISE:

Consider the one-dimensional equation,

$$w(x,t) = \frac{1}{\sqrt{2\pi\epsilon t}} \int_{-\infty}^{\infty} \bar{\rho}(\xi) e^{-(x-bt-\xi)^2/2\epsilon t} d\xi .$$

If $\bar{\rho}(\xi) = 1/2\ell$, use integration by parts to derive an expansion for $w(x,t)$.

SOLUTION:

Substituting in for $\bar{\rho}(\xi)$ gives

$$w(x,t) = \frac{1}{\sqrt{2\pi\epsilon t}} \frac{1}{2\ell} \int_{-\ell}^{\ell} e^{-(x-bt-\xi)^2/2\epsilon t} d\xi$$

$$= \frac{1}{\sqrt{2\pi\epsilon t}} \left\{ \int_{-\infty}^{\ell} e^{-(x-bt-\xi)^2/2\epsilon t} d\xi - \int_{-\infty}^{-\ell} e^{-(x-bt-\xi)^2/2\epsilon t} d\xi \right\}$$

$$= \frac{1}{\sqrt{2\pi\epsilon t}} \left\{ \int_{-\infty}^{\infty} e^{-(x-bt-\xi)^2/2\epsilon t} d\xi \right.$$

$$\left. - \int_{\ell}^{\infty} e^{-(x-bt-\xi)^2/2\epsilon t} d\xi - \int_{-\infty}^{-\ell} e^{-(x-bt-\xi)^2/2\epsilon t} d\xi \right\}$$

$$\begin{aligned}
&= \frac{1}{\sqrt{2\pi et}} \left\{ \sqrt{2\pi et} - \int_l^\infty e^{-(x-bt-\xi)^2/2et} d\xi - \int_{-\infty}^l e^{-(x-bt-\xi)^2/2et} d\xi \right\} \\
&= \frac{1}{\sqrt{2\pi et}} \left\{ \sqrt{2\pi et} - \int_l^\infty +\frac{\epsilon t}{(x-bt-\xi)} \frac{d}{d\xi} \left[e^{-(x-bt-\xi)^2/2et} \right] d\xi \right. \\
&\quad \left. - \int_{-\infty}^l +\frac{\epsilon t}{(x-bt-\xi)} \frac{d}{d\xi} \left[e^{-(x-bt-\xi)^2/2et} \right] d\xi \right\}.
\end{aligned}$$

Now integrate by parts

$$\begin{aligned}
w(x,t) &= \frac{1}{\sqrt{2\pi et}} \left\{ \sqrt{2\pi et} - \frac{\epsilon t}{(x-bt-l)} e^{-(x-bt-l)^2/2et} \right. \\
&\quad + \int_{-\infty}^l \frac{\epsilon t}{(x-bt-\xi)^2} e^{-(x-bt-\xi)^2/2et} d\xi - \frac{\epsilon t}{(x-bt+l)} e^{-(x-bt+l)^2/2et} \\
&\quad \left. + \int_{-\infty}^l \frac{\epsilon t}{(x-bt-\xi)^2} e^{-(x-bt-\xi)^2/2et} d\xi \right\}.
\end{aligned}$$

Now repeat the process.

EXERCISE:

For solution of the search equation taking the form,

$$f(x,t) = g(x,t;\epsilon) \exp[-\phi(x,t;\epsilon)] ,$$

pick $g(x,t;\epsilon)$, $\phi(x,t;\epsilon)$ and find the b^i necessary to satisfy the search equation.

SOLUTION:

This is an audience participation problem. It would defeat the purpose to give a worked example.

EXERCISE:

Using the ray ansatz,

$$f(x, t; z) = \left\{ \sum_{k=0}^{\infty} \epsilon^k g_k(x, t; z) \right\} e^{-\phi(x, t)/\epsilon}.$$

The equation for $\phi(x, t)$ is obtained by setting the coefficient of $\frac{1}{\epsilon} e^{-\phi/\epsilon}$ obtained from the search equation equal to zero. Show that this equation is

$$\frac{\partial \phi}{\partial t} + \sum_i b_i \frac{\partial \phi}{\partial x^i} + \frac{1}{2} \sum_{i,j} a^{ij} \left(\frac{\partial \phi}{\partial x^i} \right) \left(\frac{\partial \phi}{\partial x^j} \right) = 0 .$$

SOLUTION:

The search equation,

$$\frac{\partial f}{\partial t} = -\psi f - \sum_i \frac{\partial}{\partial x_i} (b_i f) + \frac{\epsilon}{2} \sum_{i,j} \frac{\partial^2 (a_{ij} f)}{\partial x_i \partial x_j} ,$$

is written equivalently as

$$\begin{aligned} \frac{\partial f}{\partial t} &= \frac{\epsilon}{2} \sum_{i,j} a_{ij} \frac{\partial^2 f}{\partial x_i \partial x_j} - \sum_i \left\{ b_i - \epsilon \sum_j \left(\frac{\partial a_{ij}}{\partial x_j} \right) \right\} \left(\frac{\partial f}{\partial x_i} \right) \\ &\quad - \left\{ \psi + \sum_i \left[\left(\frac{\partial b_i}{\partial x_i} \right) - \frac{\epsilon}{2} \sum_j \left(\frac{\partial^2 a_{ij}}{\partial x_i \partial x_j} \right) \right] \right\} f . \end{aligned}$$

$$\text{For } f(x, t; z) = \left\{ \sum_{k=0}^{\infty} \epsilon^k g_k(x, t; z) \right\} e^{-\phi(x, t)/\epsilon},$$

$$\frac{\partial f}{\partial t} = \left\{ \sum_{k=0}^{\infty} \left[\epsilon^k \frac{\partial g_k}{\partial t} - \epsilon^{k-1} \left(\frac{\partial \phi}{\partial t} \right) g_k \right] \right\} e^{-\phi/\epsilon},$$

$$\frac{\partial f}{\partial x_1} = \left\{ \sum_{k=0}^{\infty} \left[\epsilon^k \left(\frac{\partial g_k}{\partial x_1} \right) - \epsilon^{k-1} \left(\frac{\partial \phi}{\partial x_1} \right) g_k \right] \right\} e^{-\phi/\epsilon},$$

and

$$\frac{\partial^2 f}{\partial x_1 \partial x_j} = \left[\sum_{k=0}^{\infty} \left\{ \epsilon^k \left(\frac{\partial^2 g_k}{\partial x_1 \partial x_j} \right) - \epsilon^{k-1} \left[2 \left(\frac{\partial g_k}{\partial x_1} \right) \left(\frac{\partial \phi}{\partial x_j} \right) + g_k \left(\frac{\partial^2 \phi}{\partial x_1 \partial x_j} \right) \right] \right. \right. \\ \left. \left. + \epsilon^{k-2} g_k \left(\frac{\partial \phi}{\partial x_1} \right) \left(\frac{\partial \phi}{\partial x_j} \right) \right\} \right] e^{-\phi/\epsilon},$$

and the search equation becomes

$$\left\{ \sum_{k=0}^{\infty} \left[\epsilon^k \left(\frac{\partial g_k}{\partial t} \right) - \epsilon^{k-1} \left(\frac{\partial \phi}{\partial t} \right) g_k \right] \right\} e^{-\phi/\epsilon} \\ = \left[\sum_{i,j} \frac{a_{ij}}{2} \sum_{k=0}^{\infty} \left\{ \epsilon^{k+1} \left(\frac{\partial^2 g_k}{\partial x_1 \partial x_j} \right) - \epsilon^k \left[2 \left(\frac{\partial g_k}{\partial x_1} \right) \left(\frac{\partial \phi}{\partial x_j} \right) + g_k \left(\frac{\partial^2 \phi}{\partial x_1 \partial x_j} \right) \right] \right. \right. \\ \left. \left. + \epsilon^{k-1} g_k \left(\frac{\partial \phi}{\partial x_1} \right) \left(\frac{\partial \phi}{\partial x_j} \right) \right\} \right] e^{-\phi/\epsilon} \\ - \left\{ \sum_i \left[b^i - \epsilon \sum_j \left(\frac{\partial a_{ij}}{\partial x_j} \right) \right] \sum_{k=0}^{\infty} \left[\epsilon^k \left(\frac{\partial g_k}{\partial x_1} \right) - \epsilon^{k-1} \left(\frac{\partial \phi}{\partial x_1} \right) g_k \right] \right\} e^{-\phi/\epsilon} \\ - \left[\left[\psi + \sum_i \left(\frac{\partial b_i}{\partial x_1} \right) - \frac{\epsilon}{2} \sum_j \left(\frac{\partial^2 a_{ij}}{\partial x_1 \partial x_j} \right) \right] \sum_{k=0}^{\infty} \epsilon^k g_k \right] e^{-\phi/\epsilon}.$$

One now equates terms with the same power of ϵ . The equation for $\phi(x,t)$ is obtained by looking at terms involving $1/\epsilon$:

$$-\left(\frac{\partial \phi}{\partial t}\right) g_0 \frac{1}{\epsilon} e^{-\phi/\epsilon} = \sum_{i,j} \frac{a_{ij}}{2} g_0 \left(\frac{\partial \phi}{\partial x_i}\right) \left(\frac{\partial \phi}{\partial x_j}\right) \frac{1}{\epsilon} e^{-\phi/\epsilon} \\ + \sum_i b_i \left(\frac{\partial \phi}{\partial x_i}\right) g_0 \frac{1}{\epsilon} e^{-\phi/\epsilon} .$$

This yields

$$\frac{\partial \phi}{\partial t} + \frac{1}{2} \sum_{i,j} a_{ij} \left(\frac{\partial \phi}{\partial x_i}\right) \left(\frac{\partial \phi}{\partial x_j}\right) + \sum_i b_i \left(\frac{\partial \phi}{\partial x_i}\right) = 0 ,$$

the desired result.

EXERCISE:

What equation does $g_0(x,t,z)$ satisfy?

SOLUTION:

This is obtained by looking at the terms involving ϵ^0 :

$$\left[\left(\frac{\partial g_0}{\partial t} \right) - \left(\frac{\partial \phi}{\partial t} \right) g_1 \right] e^{-\phi/\epsilon} = \sum_{i,j} \frac{a_{ij}}{2} \left\{ g_1 \left(\frac{\partial \phi}{\partial x_i} \right) \frac{\partial \phi}{\partial x_j} \right. \\ \left. - \left[2 \left(\frac{\partial g_0}{\partial x_i} \right) \left(\frac{\partial \phi}{\partial x_j} \right) + g_0 \left(\frac{\partial^2 \phi}{\partial x_i \partial x_j} \right) \right] \right\} e^{-\phi/\epsilon}$$

$$\begin{aligned}
& - \sum_i b_i \left[\left(\frac{\partial g_0}{\partial x_i} \right) - \left(\frac{\partial \phi}{\partial x_i} \right) g_1 \right] e^{-\phi/c} \\
& - \sum_{i,j} \left[\left(\frac{\partial a_{ij}}{\partial x_i} \right) \left(\frac{\partial \phi}{\partial x_j} \right) g_0 \right] e^{-\phi/c} \\
& - \left[\psi + \sum_i \left(\frac{\partial b_i}{\partial x_i} \right) \right] g_0 e^{-\phi/c} .
\end{aligned}$$

Upon rearranging terms, one obtains

$$\begin{aligned}
& \frac{\partial g_0}{\partial t} + \sum_{i,j} \left\{ \frac{a_{ij}}{2} \left[2 \left(\frac{\partial g_0}{\partial x_i} \right) \left(\frac{\partial \phi}{\partial x_j} \right) + g_0 \left(\frac{\partial^2 \phi}{\partial x_i \partial x_j} \right) \right] + \left[\left(\frac{\partial a_{ij}}{\partial x_i} \right) \left(\frac{\partial \phi}{\partial x_j} \right) g_0 \right] \right\} \\
& + \sum_i \left\{ b_i \left(\frac{\partial g_0}{\partial x_i} \right) + g_0 \left(\frac{\partial b_i}{\partial x_i} \right) \right\} + \psi g_0 = 0 ,
\end{aligned}$$

since $g_1 \left\{ \frac{\partial \phi}{\partial t} + \sum_{i,j} \frac{a_{ij}}{2} \left(\frac{\partial \phi}{\partial x_i} \right) \left(\frac{\partial \phi}{\partial x_j} \right) + \sum_i b_i \left(\frac{\partial \phi}{\partial x_i} \right) \right\}$ vanishes.

(The term in braces is the left hand side of the equation just obtained for ϕ .)

EXERCISE:

Assume $a_{ij} = \delta_{ij}$ and $b_i(x,t) = b_i$, a constant. Write and solve the ray equations.

SOLUTION:

The ray equations are the following:

$$\frac{dx_i}{dt} = \frac{\partial H}{\partial p_i} \quad \text{with } x_i(0) = x_{i0}$$

and

$$\frac{dp_i}{dt} = -\frac{\partial H}{\partial x_i} \quad \text{with } p_i(0) = p_{i0},$$

where

$$H(x,p) = \sum_i b_i p_i + \frac{1}{2} \sum_{i,j} a_{ij} p_i p_j.$$

For $a_{ij} = \delta_{ij}$,

$$H(x,p) = \sum_i (b_i p_i + \frac{1}{2} p_i^2).$$

Now

$$\frac{dx_i}{dt} = \frac{\partial H}{\partial p_i} = b_i + p_i$$

and

$$\frac{dp_i}{dt} = -\frac{\partial H}{\partial x_i} = 0.$$

The second of these yields the solution, $p_i(t) = p_{i0}$.

Then, integration of the first yields

EXERCISE:

$$x_i(t) - x_{i0} = \int_0^t (b_i + p_i) dt' = (b_i + p_{i0}) t$$

and consequently,

SOLUTION

$$x_i(t) = x_{i0} + (b_i + p_{i0}) t$$

These rays are simply straight lines.

$$\frac{dx}{dt} = b_i + p_i$$

$$x = x_{i0} + (b_i + p_{i0}) t$$

$$x = x_{i0} + \frac{1}{2} (b_i + p_{i0}) t^2$$

$$(b_i + p_{i0})^2 = 1.1 \times 10^4$$

$$x = x_{i0} + \frac{1}{2} (1.1 \times 10^4) t^2$$

EXERCISE:

Assume $a_{ij} = \delta_{ij}$ and that $b_i(x, t)$ is a function of t only,
i.e., $b_i(x, t) = v_i(t)$. Write the ray equation.

SOLUTION:

$$H(x, p) = \sum_i b_i p_i + \frac{1}{2} \sum_{i,j} a_{ij} p_i p_j .$$

For $a_{ij} = \delta_{ij}$ and $b_i(x, t) = v_i(t)$,

$$H(x, p) = \sum_i \left[v_i(t) p_i + \frac{1}{2} p_i^2 \right] .$$

RAY
EQUATIONS

$$\begin{cases} \frac{dx_i}{dt} = \frac{\partial H}{\partial p_i} = p_i(t) + v_i(t) & x_i(0) = x_{i0} \\ \frac{dp_i}{dt} = -\frac{\partial H}{\partial x_i} = 0 & p_i(t) = p_i(0) = p_{i0} \end{cases} .$$

Integrating the equation for x^i ,

$$x_i(t) - x_i(0) = \int_0^t [p_i(t') + v_i(t')] dt' ,$$

and consequently,

$$x_i(t) = x_{i0} + p_{i0} t + \int_0^t v_i(t') dt'$$

EXERCISE:

What are the ray equations for general $a_{ij}(x,t)$ and $b_i(x,t)$?

SOLUTION:

For $a_{ij}(x,t)$ and $b_i(x,t)$ generally,

$$H(x,p) = \sum_i b_i(x,t)p_i(t) + \frac{1}{2} \sum_{i,j} a_{ij}(x,t)p_i(t)p_j(t) ,$$

and consequently,

$$\frac{\partial H}{\partial p_i} = b_i(x,t) + \sum_j a_{ij}(x,t)p_j(t)$$

and

$$\frac{\partial H}{\partial x_i} = \sum_i p_i(t) \left(\frac{\partial b_i}{\partial x_i} \right) + \frac{1}{2} \sum_{i,j} p_i(t)p_j(t) \left(\frac{\partial a_{ij}}{\partial x_i} \right) .$$

∴ The ray equations can be written

$$\frac{dx_i}{dt} = \frac{\partial H}{\partial p_i} = b_i(x,t) + \sum_j a_{ij}(x,t)p_j(t) ; \quad x_i(0) = x_{i0}$$

and

$$\frac{dp_i}{dt} = - \frac{\partial H}{\partial x_i} = - \sum_i p_i(t) \left(\frac{\partial b_i}{\partial x_i} \right) - \frac{1}{2} \sum_{i,j} p_i(t)p_j(t) \left(\frac{\partial a_{ij}}{\partial x_i} \right) .$$

EXERCISE:

$$\text{Suppose } \rho_0(x) = \frac{1}{2\pi\sigma^2} e^{-[(x_1)^2 + (x_2)^2]/2\sigma^2}$$

What are ϕ and h_k ($k = 0, 1, 2, \dots$) if ρ_0 is written in the form

$$\rho_0(x) = e^{-\phi(x)/\epsilon} \sum_{k=0}^{\infty} h_k(x) \epsilon^k ?$$

SOLUTION:

First, equate exponentials:

$$-\phi(x)/\epsilon = -[(x_1)^2 + (x_2)^2]/2\sigma^2 .$$

This yields

$$\phi(x) = \frac{\epsilon [(x_1)^2 + (x_2)^2]}{2\sigma^2} .$$

Next, equate the remaining terms:

$$\sum_{k=0}^{\infty} h_k(x) \epsilon^k = \frac{1}{2\pi\sigma^2} .$$

If the $h_k(x)$ are assumed to be zeroth order in ϵ , the above equation gives

$$h_0(x) = \frac{1}{2\pi\sigma^2} \quad \text{and} \quad h_k(x) = 0 \quad \text{for } k \neq 0 .$$

EXERCISE:

The Lagrangian $L\left(\mathbf{x}, \frac{d\mathbf{x}}{dt}\right)$ is defined such that

$$L\left(\mathbf{x}, \frac{d\mathbf{x}}{dt}\right) + H(\mathbf{x}, \mathbf{p}) = \sum_i \frac{dx_i}{dt} \cdot p_i .$$

Show that

$$L\left(\mathbf{x}, \frac{d\mathbf{x}}{dt}\right) = \frac{1}{2} \sum_{i,j} \left(\frac{dx_i}{dt} - b_i \right) \left(a_{ij} \right)^{-1} \left(\frac{dx_j}{dt} - b_j \right) ,$$

where $\left(a_{ij}\right)^{-1}$ is the i,j^{th} element of the inverse of the matrix $[a_{ij}]$.

SOLUTION:

$$H(\mathbf{x}, \mathbf{p}) = \sum_i b_i p_i + \frac{1}{2} \sum_{i,j} a_{ij} p_i p_j$$

and

$$\frac{dx_i}{dt} = \frac{\partial H}{\partial p_i} = b_i + \sum_j a_{ij} p_j .$$

Now,

$$\frac{dx_i}{dt} - b_i = \sum_j a_{ij} p_j$$

can be written as

$$\left(\frac{dx}{dt} - \underline{b} \right)_i = \left([a_{ij}] \cdot \underline{p} \right)_i .$$

where $\left(\frac{dx}{dt} - b \right)_i$ is the i^{th} row of the $n \times 1$ column matrix,

the elements of which are the components of the vector $\left(\frac{d}{dt} x - b \right)$.

$[a_{ij}] \cdot p$ is the product of the $n \times m$ matrix with elements

a_{ij} and the column matrix ($n \times 1$) whose elements are the components of the vector p .

Since

$$\left(\frac{dx}{dt} - b \right) = [a_{ij}] \cdot p ,$$

$$p = [a_{ij}]^{-1} \cdot \left(\frac{dx}{dt} - b \right) ,$$

where $[a_{ij}]^{-1}$ is the inverse of the matrix $[a_{ij}]$.

Now

$$p_k = \left([a_{ij}]^{-1} \cdot \left(\frac{dx}{dt} - b \right) \right)_k$$

$$= \sum_l \left([a_{ij}]^{-1} \right)_{kl} \left(\frac{dx_l}{dt} - b_l \right) ,$$

where $\left([a_{ij}]^{-1} \right)_{kl}$ is the k, l^{th} element of the matrix that is

the inverse of the matrix $[a_{ij}]$.

Turning to the Lagrangian,

$$\begin{aligned}
 L\left(\underline{x}, \frac{d\underline{x}}{dt}\right) &= \sum_i \frac{d\underline{x}_i}{dt} \cdot p_i - H(\underline{x}, \underline{p}) \\
 &= \underbrace{\sum_i b_i p_i}_{\sum_i \frac{d\underline{x}_i}{dt} \cdot p_i} + \underbrace{\sum_{i,j} a_{ij} p_j p_i}_{\sum_i b_i p_i + \frac{1}{2} \sum_{i,j} a_{ij} p_i p_j} - \left[\underbrace{\sum_i b_i p_i}_{H(\underline{x}, \underline{p})} + \frac{1}{2} \sum_{i,j} a_{ij} p_i p_j \right] \\
 &= \frac{1}{2} \sum_{i,j} a_{ij} p_i p_j \\
 &= \frac{1}{2} (\underline{p})^{\text{Tr}} \cdot [\underline{a}_{ij}] \cdot (\underline{p})
 \end{aligned}$$

where $(\underline{p})^{\text{Tr}}$ is the $1 \times n$ row matrix obtained by taking the transpose of the $n \times 1$ column matrix (\underline{p}) .

$$\begin{aligned}
 (\underline{p})^{\text{Tr}} &= \left(\frac{d\underline{x}}{dt} - \underline{b} \right)^{\text{Tr}} \left([\underline{a}_{ij}]^{-1} \right)^{\text{Tr}} \\
 &= \left(\frac{d\underline{x}}{dt} - \underline{b} \right)^{\text{Tr}} \left([\underline{a}_{ij}^{\text{Tr}}]^{-1} \right)^{-1} \\
 &= \left(\frac{d\underline{x}}{dt} - \underline{b} \right)^{\text{Tr}} [\underline{a}_{ij}^{\text{Tr}}]^{-1}
 \end{aligned}$$

Now,

$$\begin{aligned} (\underline{\underline{p}})^T \underline{\underline{a}_{ij}} \cdot (\underline{\underline{p}}) &= \left(\frac{dx}{dt} - b \right)^T \underline{\underline{a}_{ij}}^{-1} \underline{\underline{a}_{ij}} \underline{\underline{a}_{ij}}^{-1} \left(\frac{dx}{dt} - b \right) \\ &= \left(\frac{dx}{dt} - b \right)^T \underline{\underline{a}_{ij}}^{-1} \left(\frac{dx}{dt} - b \right) \\ &= \sum_{kl} \left(\frac{dx_k}{dt} - b_k \right) \left(\underline{\underline{a}_{ij}}^{-1} \right)_{kl} \left(\frac{dx_l}{dt} - b_l \right) \\ &= \sum_{kl} \left(\frac{dx_k}{dt} - b_k \right) \left(a_{ij} \right)^{-1} \left(\frac{dx_l}{dt} - b_l \right) , \end{aligned}$$

if one defines $\left(a_{ij} \right)^{-1} = \left(\underline{\underline{a}_{ij}}^{-1} \right)_{kl}$ to be the k, l^{th} element

of the inverse of the matrix with elements a_{ij} .

Finally,

$$\begin{aligned} L\left(x, \frac{dx}{dt}\right) &= \frac{1}{2} \sum_{i,j} a_{ij} p_i p_j = \frac{1}{2} (\underline{\underline{p}})^T \cdot \underline{\underline{a}_{ij}} \cdot (\underline{\underline{p}}) \\ &= \frac{1}{2} \sum_{i,j} \left(\frac{dx_i}{dt} - b_i \right) \left(a_{ij} \right)^{-1} \left(\frac{dx_j}{dt} - b_j \right) , \end{aligned}$$

which was to be shown.

CNA Professional Papers - 1973 to Present*

- PP 103** Friedheim, Robert L., "Political Aspects of Ocean Security," 48 pp., Feb 1973, published in Who Presents the Ocean, John Lawrence Margrove (ed.) (St. Paul: West Publ. Co., 1974), published by the American Society of International Law) AD 757 236
- PP 104** Schles, Jack M., "A Review of James Cable, Gunboat Diplomacy: Political Applications of Limited Naval Forces," 5 pp., Feb 1973, (Reviewed in the American Political Science Review, Vol. LXVI, Dec 1972)
- PP 105** Cern, Robert J. and Phillips, Gary R., "On Optimal Correction of Gunfire Errors," 22 pp., Mar 1973, AD 751 674
- PP 106** Stoloff, Peter H., "User's Guide for Generalized Factor Analysis Program (PACTAN)," 26 pp., Feb 1973, (Includes an addendum published Aug 1974) AD 758 824
- PP 107** Stoloff, Peter H., "Relating Factor Analytically Derived Measures to Exogenous Variables," 17 pp., Mar 1973, AD 758 820
- PP 108** McConnell, James M. and Kelly, Anne M., "Super-power Naval Diplomacy in the Indo-Pakistani Crisis," 14 pp., 8 Feb 1973, (Published, with revisions, in Survival, Nov/Dec 1973) AD 751 678
- PP 109** Berghoef, Fred G., "Societies-A Framework for the Study of Trend," 8 pp., Dec 1973, (Published in Review of Income and Wealth, Series 19, No. 4, Dec 1973)
- PP 110** Auguste, Joseph, "A Critique of Cost Analysis," 9 pp., Jul 1973, AD 756 576
- PP 111** Hendrik, Robert W., "The USSR's 'Blue Belt of Defense' Concept: A Unified Military Plan for Defense Against Submarine Nuclear Attack by Strike Carriers and Polaris/Poseidon SSBNs," 18 pp., May 1973, AD 756 375
- PP 112** Glimborg, Lawrence H., "ELF Atmosphere Noise Level Statistics for Project SANGUINE," 29 pp., Apr 1974, AD 758 809
- PP 113** Glimborg, Lawrence H., "Propagation Anomalies During Project SANGUINE Experiments," 5 pp., Apr 1974, AD 754 808
- PP 114** Meloney, Arthur P., "Job Satisfaction and Job Turnover," 41 pp., Jul 1973, AD 758 410
- PP 115** Silverman, Lester P., "The Determinants of Emergency and Elective Admissions to Hospitals," 145 pp., 18 Jul 1973, AD 758 377
- PP 116** Rohm, Allan S., "An Assessment of Military Operations Research in the USSR," 19 pp., Sep 1973, (Reprinted from Proceedings, 3rd Military Operations Research Symposium (U), Secret Dec 1972) AD 770 116
- PP 117** McWhirter, Peter B. and Ratliff, H. Donald,* "Defending a Logistics System Under Mining Attack," 24 pp., Aug 1973 (to be submitted for publication in Naval Research Logistics Quarterly), presented at 44th National Meeting, Operations Research Society of America, November 1973, AD 830 484
*University of Florida.
**Research supported in part under Office of Naval Research Contract N00014-68-0273-0017
- PP 118** Barfoot, C. Bernard, "Market Duds," 18 pp., Apr 1973, (Reprinted from Operations Research, Vol. 22, No. 2, Mar-Apr 1974)
- PP 119** Stoloff, Peter and Lockman, Robert F., "Development of Navy Human Relations Questionnaire," 8 pp., May 1974, (Published in American Psychological Association Proceedings, 81st Annual Convention, 1973) AD 758 240
- PP 120** Smith, Michael W. and Schwipper, Ronald A., "Econometric Analysis of the Intraday Dispersion of Criminal Activity," 30 pp., Jun 1974, (Presented at the Econometrics Society Meetings, 30 Dec 1973) AD 760 520
*Economics, North Carolina State University.
- PP 121** Devine, Eugene J., "Procurement and Retention of Navy Physicians," 21 pp., Jun 1974, (Presented at the 49th Annual Conference, Western Economic Association, Las Vegas, Nev., 10 Jun 1974) AD 760 539
- PP 122** Kelly, Anne M., "The Soviet Naval Presence During the Iraq-Kuwait Border Dispute: March-April 1973," 34 pp., Jun 1974, (Published in Soviet Naval Policy, ed. Michael McGwire; New York: Praeger) AD 750 888
- PP 123** Petersen, Charles G., "The Soviet Port-Closing Operation in Bangladesh, March 1972-December 1973," 38 pp., Jun 1974, (Published in Michael McGwire, et al. (eds) Soviet Naval Policy: Objectives and Constraints, (New York: Praeger Publishers, 1974) AD 780 540
- PP 124** Friedheim, Robert L. and John, Mary E., "Anticipating Soviet Behavior at the Third U.N. Law of the Sea Conference: USSR Positions and Dilemmas," 27 pp., 10 Apr 1974, (Published in Soviet Naval Policy, ed. Michael McGwire; New York: Praeger) AD 753 701
- PP 125** Weinland, Robert G., "Soviet Naval Operations-Ten Years of Change," 17 pp., Aug 1974, (Published in Soviet Naval Policy, ed. Michael McGwire; New York: Praeger) AD 753 902
- PP 126** -- Classified.
- PP 127** Dragovich, George S., "The Soviet Union's Quest for Access to Naval Facilities in Egypt Prior to the June War of 1967," 64 pp., Jul 1974, AD 756 316
- PP 128** Stoloff, Peter and Lockman, Robert F., "Evaluation of Naval Officer Performance," 11 pp., (Presented at the 83rd Annual Convention of the American Psychological Association, 1974) Aug 1974, AD 754 012
- PP 129** Holen, Arlene and Horowitz, Stanley, "Partial Unemployment Insurance Benefits and the Extent of Partial Unemployment," 4 pp., Aug 1974, (Published in the Journal of Human Resources, Vol. IX, No. 3, Summer 1974) AD 754 010
- PP 130** Dimukos, Bradford, "Roles and Missions of Soviet Naval General Purpose Forces in Wartime Pro-SBNN Operation," 20 pp., Aug 1974, AD 758 320
- PP 131** Weinland, Robert G., "Analysis of Gorbachev's Naval in War and Peace," 48 pp., Aug 1974, (Published in Soviet Naval Policy, ed. Michael McGwire; New York: Praeger) AD 756 318
- PP 132** Kleinman, Samuel D., "Racial Differences in Hours Worked in the Market: A Preliminary Report," 27 pp., Feb 1975, (Paper read on 26 Oct 1974 at Eastern Economic Association Convention in Albany, N.Y.) AD A 006 817
- PP 133** Squires, Michael L., "A Stochastic Model of Regime Change in Latin America," 48 pp., Feb 1975, AD A 007 812
- PP 134** Root, R. M. and Cunniff, P. F.,* "A Study of the Shock Spectrum of a Two-Degree-of-Freedom Nonlinear Vibratory System," 39 pp., Dec 1975, (Published in the condensed version of The Journal of the Acoustical Society, Vol. 60, No. 6, Dec 1976, pp. 1314)
*Department of Mechanical Engineering, University of Maryland.
- PP 135** Goudreau, Kenneth A.; Kusmark, Richard A.; Wiedemann, Karen, "Analysis of Closure Alternatives for Naval Stations and Naval Air Stations," 47 pp., 3 Jun 1975, (Reprinted from "Hearing before the Subcommittee on Military Construction of the Committee on Armed Services," U.S. Senate, 93rd Congress, 1st Session, Part 2, 22 Jun 1973)
- PP 136** Stahlings, William, "Cybernetics and Behavior Therapy," 12 pp., Jun 1975
- PP 137** Peterson, Charles G., "The Soviet Union and the Reopening of the Suez Canal: Miniclearing Operations in the Gulf of Suez," 30 pp., Aug 1975, AD A 018 378

*CNA Professional Papers with an AD number may be obtained from the National Technical Information Service, U.S. Department of Commerce, Springfield, Virginia 22181. Other papers are available from the author at the Center for Naval Analyses, 2000 North Beauregard Street, Alexandria, Virginia 22311.

- PP 138
Balling, William, "BRIDGE: An Interactive Dialogue-Generation Facility," 6 pp., Aug 1976 (Reprinted from IEEE Transactions on Systems, Man, and Cybernetics, Vol. S No. 3, May 1978)
- PP 139
Morgan, William F., Jr., "Beyond Politics and Pebbles in Poetry to Positive Economics," 14 pp., (Presented at Southern Economic Association Meetings November, 1974) Aug 1976, AD A 018 283
- PP 140
Mahoney, Robert and Druckman, Daniel, "Simulation, Experimentation, and Context," 38 pp., 1 Sep 1976, (Published in Simulation & Games, Vol. 6, No. 3, Sep 1976)
"Mathematics, Inc."
- PP 141
Mizrahi, Maurice M., "Generalized Hermite Polynomials," 8 pp., Feb 1976 (Reprinted from the Journal of Computational and Applied Mathematics, Vol. 1, No. 4 (1976), 273-277).
"Research supported by the National Science Foundation"
- PP 142
Lockman, Robert F., John, Christopher, and Shugart, William F. II, "Models for Estimating Premature Losses and Recruiting District Performance," 36 pp., Dec 1976 (Presented at the RAND Conference on Defense Manpower, Feb 1976; to be published in the conference proceedings) AD A 020 442
- PP 143
Horowitz, Stanley and Sherman, Allan (LCDR, USN), "Maintenance Personnel Effectiveness in the Navy," 33 pp., Jan 1976 (Presented at the RAND Conference on Defense Manpower, Feb 1976; to be published in the conference proceedings) AD A 021 561
- PP 144
Dureh, William J., "The Navy of the Republic of China - History, Problems, and Prospects," 88 pp., Aug 1976 (To be published in "A Guide to Asiatic Fleets," ed. by Barry M. Blechman; Naval Institute Press) AD A 020 460
- PP 145
Kelly, Anne M., "Port Visits and the 'Internationalist Mission' of the Soviet Navy," 26 pp., Apr 1976 AD A 023 430
- PP 146
Palmer, Vernon E., "Alternatives for Increasing Access to Scientific Journals," 8 pp., Apr 1976 (Presented at the 1976 IEEE Conference on Scientific Journals, Cherry Hill, N.C., Apr 28-30; published in IEEE Transactions on Professional Communication, Vol. PC-18, No. 3, Sep 1976) AD A 021 798
- PP 147
Kessler, J. Christian, "Legal Issues in Protecting Offshore Structures," 39 pp., Jun 1976 (Prepared under task order N00014-68-A-0091-0023 for ONR) AD A 028 389
- PP 148
McConnell, James M., "Military-Political Tasks of the Soviet Navy in War and Peace," 62 pp., Dec 1976 (Published in Soviet Ocean Development Study of Senate Commerce Committee October 1976) AD A 022 590
- PP 149
Squires, Michael L., "Counterforce Effectiveness: A Comparison of the Tripis 'K' Measure and a Computer Simulation," 24 pp., Mar 1976 (Presented at the International Study Association Meeting, 27 Feb 1976) AD A 022 591
- PP 150
Kelly, Anne M. and Peterkin, Charles, "Recent Changes in Soviet Naval Policy: Prospects for Arms Limitations in the Mediterranean and Indian Ocean," 28 pp., Apr 1976, AD A 023 723
- PP 151
Horowitz, Stanley A., "The Economic Consequences of Political Philosophy," 8 pp., Apr 1976 (Reprinted from Economic Inquiry, Vol. XIV, No. 1, Mar 1976)
- PP 152
Mizrahi, Maurice M., "On Path Integral Solutions of the Schrödinger Equation, Without Limiting Procedure," 10 pp., Apr 1976 (Reprinted from Journal of Mathematical Physics, Vol. 17, No. 4 (Apr 1976), 566-578).
"Research supported by the National Science Foundation"
- PP 153
Mizrahi, Maurice M., "WKB Expansions by Path Integrals, With Applications to the Anharmonic Oscillator," 137 pp., May 1976, AD A 026 440
"Research supported by the National Science Foundation"
- PP 154
Mizrahi, Maurice M., "On the Semi-Classical Expansion in Quantum Mechanics for Arbitrary Hamiltonians," 16 pp., May 1976 (Published in Journal of Mathematical Physics, Vol. 18, No. 4, p. 788, Apr 1977), AD A 026 441
- PP 155
Squires, Michael L., "Soviet Foreign Policy and Third World Nations," 20 pp., Jun 1976 (Prepared for presentation at the Midwest Political Science Association meetings, Apr 30, 1976) AD A 028 388
- PP 156
Balling, William, "Approaches to Chinese Character Recognition," 12 pp., Jun 1976 (Reprinted from Pattern Recognition (Pergamon Press), Vol. 8, pp. 87-98, 1976) AD A 028 692
- PP 157
Morgan, William F., "Unemployment and the Pentagon Budget: Is There Anything in the Empty Pork Barrel?" 20 pp., Aug 1976 AD A 030 488
- PP 158
Haskell, LCDR, Richard D. (USN), "Experimental Validation of Probability Predictions," 28 pp., Aug 1976 (Presented at the Military Operations Research Society Meeting, Fall 1976) AD A 020 458
- PP 159
McConnell, James M., "The Gorshkov Articles, The New Gorshkov Book and Their Relation to Policy," 83 pp., Jul 1976 (Published in Soviet Naval Influence: Domestic and Foreign Dimensions, ed. by M. Meelwiss and J. McDonald; New York: Praeger, 1977) AD A 020 227
- PP 160
Wilson, Desmond P., Jr., "The U.S. Sixth Fleet and the Conventional Defense of Europe," 60 pp., Sep 1976 (Submitted for publication in Adelphi Papers, 118 S., London) AD A 030 487
- PP 161
Meltch, Michael E. and Peet, Vice Adm. Roy (USN, Retired), "Fleet Commanders: Allot or Author?" 9 pp., Aug 1976 (Reprinted from U.S. Naval Institute Proceedings, Jun 1976) AD A 030 456
- PP 162
Friedheim, Robert L., "Parliamentary Diplomacy," 108 pp., Sep 1976 AD A 033 306
- PP 163
Lockman, Robert F., "A Model for Predicting Re-Entry Losses," 8 pp., Sep 1976 (Presented at the 84th annual convention of the American Psychological Association, Washington, D.C., 4 Sep 1976) AD A 030 489
- PP 164
Mahoney, Robert F., Jr., "An Assessment of Public and Elite Perceptions in France, The United Kingdom, and the Federal Republic of Germany, 31 pp., Feb 1977 (Presented at Conference "Perception of the U.S. - Soviet Balance and the Political Uses of Military Power" sponsored by Director, Advanced Research Projects Agency, April 1976) AD A 026 699
- PP 165
Jondrow, James M. "Effects of Trade Restrictions on Imports of Steel," 67 pp., November 1976, (Delivered at ILAB Conference in Dec 1976)
- PP 166
Feldman, Paul, "Impediments to the Implementation of Desirable Changes in the Regulation of Urban Public Transportation," 12 pp., Oct 1976, AD A 033 322
- PP 167 - Revised
Feldman, Paul, "Why It's Difficult to Change Regulation," Oct 1976
- PP 168
Kleinman, Samuel, "ROTC Service Commitments: a Comment," 4 pp., Nov 1976, (To be published in Public Choice, Vol. XXIV, Fall 1976) AD A 032 308
- PP 169
Lockman, Robert F., "Revalidation of CNA Support Personnel Selection Measures," 30 pp., Nov 1976
- PP 170
Jacobsen, Louis B., "Earnings Losses of Workers Displaced from Manufacturing Industries," 38 pp., Nov 1976, (Delivered at ILAB Conference in Dec 1976) AD A 030 608
- PP 171
Raiton, James M., "A Diffusion Model for Gap Red LED Degradation," 10 pp., Nov 1976, (Published in Journal of Applied Physics, Vol. 47, pp. 4818-4827, Oct 1976)

- PP 172
Clossen, Kathleen P., "Unemployment Insurance and the Length of Unemployment," Dec 1976, (Presented at the University of Rochester Labor Workshop on 18 Nov 1976)
- PP 173
Kleiman, Samuel D., "A Note on Racial Differences in the Active-Worker/Diseconomized-Worker Controversy," 8 pp., Dec 1976, (Published in the American Economist, Vol. XX, No. 1, Spring 1976)
- PP 174
Mahoney, Robert E., Jr., "A Comparison of the Beckoning and International Incident Projects," 12 pp., Feb 1977 AD A040 206
- PP 175
Levine, Donald; Stoltz, Peter and Spratt, Neway, "Public Drug Treatment and Addict Crime," June 1976, (Published in Journal of Legal Studies, Vol. 5, No. 2)
- PP 176
Folia, Wendi, "Correlates of Retention and Promotion for USNA Graduates," 38 pp., Mar 1977, AD A000 040
- PP 177
Lockman, Robert F. and Warner, John T., "Predicting Attrition: A Test of Alternative Approaches," 33 pp., Mar 1977, (Presented at the OBD/ONR Conference on Enlisted Attrition Xerox International Training Center, Leesburg, Virginia, 47 April 1977), AD A039 047
- PP 178
Kleinman, Samuel D., "An Evaluation of Navy Unrestricted Line Officer Acquisition Programs," 23 pp., April 1977, (To be presented at the NATO Conference on Manpower Planning and Organization Design, Stresa, Italy, 20 June 1977), AD A039 048
- PP 179
Stoltz, Peter H. and Salut, Stephen J., "Vehicle: A Model for Personnel Inventory Planning Under Changing Management Policy," 14 pp., April 1977, (Presented at the NATO Conference on Manpower Planning and Organization Design, Stresa, Italy, 20 June 1977), AD A039 049
- PP 180
Horowitz, Stanley A. and Sherman, Allan, "The Characteristics of Naval Personnel and Personnel Performance," 10 pp., April 1977, (Presented at the NATO Conference on Manpower Planning and Organization Design, Stresa, Italy, 20 June 1977), AD A039 050
- PP 181
Salut, Stephen J. and Stoltz, Peter, "An Inventory Manning Model for Navy Enlisted Personnel," 38 pp., May 1977, (Prepared for presentation at the Joint National Meeting of the Operations Research Society of America and The Institute for Management Sciences, 9 May 1977, San Francisco, California), AD A048 221
- PP 182
Murray, Russell, End, "The Quest for the Perfect Study or My First 1138 Days at CNA," 67 pp., April 1977
- PP 183
Kassing, David, "Changes in Soviet Naval Forces," 23 pp., November, 1976, (Published as part of Chapter 3, "General Purpose Forces: Navy and Marine Corps," in Arms, Men, and Military Budgets, Francis P. Hooper and William Schneider, Jr. (eds.), (Crane, Russak & Company, Inc.: New York), 1977), AD A040 106
- PP 184
Lockman, Robert F., "An Overview of the OBD/ONR Conference on First Term Enlisted Attrition," 22 pp., June 1977, (Presented to the 30th MCRA Working Group on Manpower and Personnel Planning, Annapolis, Md., 28-30 June 1977), AD A048 918
- PP 185
Kassing, David, "New Technology and Naval Forces in the South Atlantic," 22 pp., (This paper was the basis for a presentation made at the Institute for Foreign Policy Analysis, Cambridge, Mass., 28 April 1977), AD A048 919
- PP 186
Mazrahi, Maurice M., "Phase Space Integrals, Without Limiting Procedure," 31 pp., May 1977, (Invited paper presented at the 1977 NATO Institute on Path Integrals and Their Application in Quantum Statistical, and Solid State Physics, Antwerp, Belgium, July 17-20, 1977) (Published in Journal of Mathematical Physics 18(1), p. 208, Jan 1977), AD A048 107
- PP 187
Coile, Russell C., "Nomography for Operations Research," 30 pp., April 1977, (Presented at the Joint National Meeting of the Operations Research Society of America and The Institute for Management Services, San Francisco, California, 8 May 1977), AD A043 820
- PP 188
Dorch, William J., "Information Processing and Outcome Forecasting for Multilateral Negotiations: Testing One Approach," 53 pp., May 1977 (Prepared for presentation at the 18th Annual Convention of the International Studies Association, Chouteau Park Plaza Hotel, St. Louis, Missouri, March 19-23, 1977), AD A048 222
- PP 189
Coile, Russell C., "Error Detection in Computerized Information Retrieval Data Bases," July, 1977, 18 pp., Presented at the Sixth Cranfield International Conference on Mechanized Information Storage and Retrieval Systems, Cranfield Institute of Technology, Cranfield, Bedford, England, 26-28 July 1977, AD A043 880
- PP 190
Mahoney, Robert E., Jr., "European Perceptions and East-West Competition," 90 pp., July 1977 (Prepared for presentation at the annual meeting of the International Studies Association, St. Louis, Mo., March, 1977), AD A043 981
- PP 191
Bawyer, Ronald, "The Independent Field Assignment: One Man's View," August 1977, 18 pp.
- PP 192
Helen, Ariane, "Effects of Unemployment Insurance Entitlement on Duration and Job Search Outcome," August 1977, 8 pp., (Reprinted from Industrial and Labor Relations Review, Vol. 30, No. 4, Jul 1977)
- PP 193
Horowitz, Stanley A., "A Model of Unemployment Insurance and the Work Tax," August 1977, 7 pp., (Reprinted from Industrial and Labor Relations Review, Vol. 30, No. 4, Jul 1977)
- PP 194
Clossen, Kathleen P., "The Effects of Unemployment Insurance on the Duration of Unemployment and Subsequent Earnings," August 1977, 7 pp., (Reprinted from Industrial and Labor Relations Review, Vol. 30, No. 4, Jul 1977)
- PP 195
Bruckling, Frank, "Unemployment Insurance Taxes and Labor Turnover: Summary of Theoretical Findings," 12 pp., (Reprinted from Industrial and Labor Relations Review, Vol. 30, No. 4, Jul 1977)
- PP 196
Rosten, J. M. and Lorimer, G. G., "Degradation of Bulk Electroluminescent Efficiency in Zn, O-Doped GaP LEDs," July 1977, 3 pp., (Reprinted from IEEE Transactions on Electron Devices, Vol. ED-24, No. 7, July 1977)
- PP 197
Wells, Anthony R., "The Centre for Naval Analyses," 14 pp., Dec 1977, AD A048 107
- PP 198
Clossen, Kathleen P., "The Distributional Effects of Unemployment Insurance," 25 pp., Sept. 1977 (Presented at a Hoover Institution Conference on Income Distribution, Oct 7-8, 1977)
- PP 199
Dorch, William J., "Revolution From A P.A.R. - The Cuban Armed Forces in Africa and the Middle East," Sep 1977, 16 pp., AD A048 206
- PP 200
Power, Bruce F., "The United States Navy," 40 pp., Dec 1977, (To be published as a chapter in The U.S. War Machine by Ballantine Books in England during 1978), AD A049 106
- PP 201
Dorch, William J., "The Cuban Military in Africa and The Middle East: From Algeria to Angola," Sep 1977, 87 pp., AD A048 678
- PP 202
Feldman, Paul, "Why Regulation Doesn't Work," (Reprinted from Technological Change and Welfare in the Regulated Industries and Review of Social Economy, Vol. XXIX, March, 1971, No. 1.) Sep 1977, 8 pp.
- PP 203
Feldman, Paul, "Efficiency, Distribution, and the Role of Government in a Market Economy," (Reprinted from The Journal of Political Economy, Vol. 78, No. 3, May/June 1971.) Sep 1977, 19 pp., AD A048 676

- PP 204
Wells, Anthony R., "The 1967 June War: Soviet Naval Diplomacy and The Sixth Fleet - A Re-appraisal," Oct 1977, 36 pp., AD A047 236
- PP 205
Colle, Russell C., "A Bibliometric Examination of the Square Root Theory of Scientific Publication Productivity," (Presented at the annual meeting of the American Society for Information Science, Chicago, Illinois, 20 September 1977.) Oct 1977, 8 pp., AD A047 237
- PP 206
McConnell, James M., "Strategy and Missions of the Soviet Navy in the Year 2000," 48 pp., Nov 1977, (Presented at a Conference on Problems of Sea Power as we Approach the 21st Century, sponsored by the American Enterprise Institute for Public Policy Research, 8 October 1977, and subsequently published in a collection of papers by the Institute), AD A047 244
- PP 207
Goldberg, Lawrence, "Cost Effectiveness of Potential Federal Policies Affecting Research & Development Expenditures in the Auto, Steel and Food Industries," 30 pp., Oct 1977, (Presented at Southern Economic Association Meetings beginning 2 November 1977)
- PP 208
Roberts, Stephen B., "The Decline of the Overseas Station Fleets: The United States Asiatic Fleet and the Shanghai Crisis, 1932," 18 pp., Nov 1977, (Reprinted from *The American Neptune*, Vol. XXXVII, No. 3, July 1977), AD A047 248
- PP 209 - Classified.
- PP 210
Kassing, David, "Protecting The Fleet," 40 pp., Dec 1977 (Presented for the American Enterprise Institute Conference on Problems of Sea Power as We Approach the 21st Century, October 6-7, 1977), AD A048 106
- PP 211
Mireshi, Maurice M., "On Approximating the Circular Coverage Function," 14 pp., Feb 1978
- PP 212
Mangel, Marc, "On Singular Characteristic Initial Value Problems with Unique Solutions," 20 pp., Jun 1978 (To be submitted for publication in Journal of Mathematical Analysis and Its Applications)
- PP 213
Mangel, Marc, "Fluctuations in Systems with Multiple Steady States. Application to Lanchester Equations," 12 pp., Feb 78, (Presented at the First Annual Workshop on the Information Linkage Between Applied Mathematics and Industry, Naval PG School, Feb 23-26, 1978)
- PP 214
Wineland, Robert G., "A Somewhat Different View of The Optimal Naval Posture," 37 pp., Jun 1978 (Presented at the 1978 Convention of the American Political Science Association (APSA)/US Panel on "Changing Strategic Requirements and Military Posture"), Chicago, Ill., September 2, 1978)
- PP 215
Colle, Russell C., "Comments on: Principle of Information Retrieval by Manfred Kuchen, 10 pp., Mar 78, (Published as a Letter to the Editor, Journal of Documentation, Vol. 31, No. 4, pages 298-301, December 1975)
- PP 216
Colle, Russell C., "Lotha's Frequency Distribution of Scientific Productivity," 18 pp., Feb 1978, (Published in the Journal of the American Society for Information Science, Vol. 29, No. 6, pp. 368-370, November 1977)
- PP 217
Colle, Russell C., "Bibliometric Studies of Scientific Productivity," 17 pp., Mar 78, (Presented at the Annual meeting of the American Society for Information Sciences held in San Francisco, California, October 1978.)
- PP 218 - Classified.
- PP 219
Huntington, R. LaVoi, "Market Analysis with Rational Expectations: Theory and Estimation," 60 pp., Apr 78 (To be submitted for publication in Journal of Econometrics)
- PP 220
Meurer, Donald E., "Diagonalization by Group Matrices," 26 pp., Apr 78
- PP 221
Weinland, Robert G., "Superpower Naval Diplomacy in the October 1973 Arab-Israeli War," 28 pp., Jun 1978
- PP 222
Mireshi, Maurice M., "Correspondence Rules and Path Integrals," 30 pp., Jun 1978 (Invited paper presented at the CNRS meeting on "Mathematical Problems in Feynman's Path Integrals," Marseille, France, May 22-26, 1978)
- PP 223
Mangel, Marc, "Stochastic Mechanics of Molecular-Ion Molecule Reactions," 21 pp., Jun 1978 (To be submitted for publication in Journal of Mathematical Physics)
- PP 224
Mangel, Marc, "Aggregation, Bifurcation, and Extinction in Unexploited Animal Populations," 48 pp., Mar 1978 (To be submitted for publication in American Naturalist)
'Portions of this work were started at the Institute of Applied Mathematics and Statistics, University of British Columbia, Vancouver, B.C., Canada'
- PP 225
Mangel, Marc, "Oscillations, Fluctuations, and the Hopf Bifurcation," 43 pp., Jun 1978
'Portions of this work were completed at the Institute of Applied Mathematics and Statistics, University of British Columbia, Vancouver, Canada.'
- PP 226
Rosten, J. M. and J. W. Mann, "Temperature and Current Dependence of Degradation in Lead-Smitting Gas LEDs," 34 pp., Jun 1978
- PP 227
Mangel, Marc, "Uniform Treatment of Fluctuations at Critical Points," 50 pp., May 1978 (To be submitted for publication in Journal of Statistical Physics)
- PP 228
Mangel, Marc, "Relaxation at Critical Points: Deterministic and Stochastic Theory," 54 pp., Jun 1978 (To be submitted for publication in Journal of Mathematical Physics)
- PP 229
Mangel, Marc, "Diffusion Theory of Reaction Rates, I: Formulation and Einstein-Smoluchowski Approximation," 50 pp., Jun 1978
- PP 230
Mangel, Marc, "Diffusion Theory of Reaction Rates, II: Ornstein-Uhlenbeck Approximation," 34 pp., Feb 1978
- PP 231
Wilson, Desmond P., Jr., "Naval Projection Forces: The Case for a Responsive NAVF," Aug 1978
- PP 232
Jacobson, Louis, "Can Policy Changes be Made Acceptable to Labor?" Aug 1978 (To be submitted for publication in Industrial and Labor Relations Review)
- PP 233
Jacobson, Louis, "An Alternative Explanation of the Cyclical Pattern of Quits," 23 pp., Sep 1978
- PP 234
Jondrow, James and Levy, Robert A., "Does Federal Expenditure Disclose State and Local Expenditure: The Case of Construction Grants," 38 pp., Oct 1978 (To be submitted for publication in Journal of Public Economics)
- PP 235
Mireshi, Maurice M., "The Semiclassical Expansion of the Anharmonic-Oscillator Propagator," 41 pp., Oct 1978 (To appear in the Journal of Mathematical Physics)
- PP 237
Meurer, Donald, "A Matrix Criterion for Normal Integral Bases," 10 pp., Jan 1979
- PP 238
Utgoff, Kathleen Classen, "Unemployment Insurance and The Employment Rate," 20 pp., Oct 1978
- PP 239
Trost, R. P. and Warner, J. T., "The Effects of Military Occupational Training on Civilian Earnings: An Income Selectivity Approach," 30 pp., Nov 1978 (To be submitted for publication in Review of Economics and Statistics)
- PP 240
Powers, Bruce, "Goals of the Center for Naval Analysts," 13 pp., Dec 1978
- PP 241
Mangel, Marc, "Fluctuations at Chemical Instabilities," 24 pp., Dec 1978 (Published in Journal of Chemical Physics, Vol. 69, No. 8, Oct 15, 1978)

- PP 242
Simpson, William R., "The Analysis of Dynamically Interactive Systems (An Contest by the Numbers)," 100 pp., Dec 1970
- PP 243
Simpson, William R., "A Probabilistic Formulation of Nuclear Dynamics as Applied to the Analysis of Operational Research Problems," 18 pp., Dec 1970 (Submitted for publication in *The Journal of Irreproducible Results*)
- PP 244
Sherman, Allan and Horowitz, Stanley A., "Maintenance Costs of Complex Equipment," 20 pp., Dec 1970
- PP 245
Simpson, William R., "The Accelerometer Methods of Observing Aircraft Performance from Flight Test Data (Dynamic Performance Testing)," 403 pp., Jun 1970
- PP 246
Thomas, James A., Jr., "The Transport Properties of Dilute Gases in Applied Fields," 182 pp., Mar 1970
- PP 247
Gleser, Kenneth S., "A Secretary Problem with a Random Number of Choices," 23 pp., Mar 1970 (Submitted for publication in *Journal of the American Statistical Association*)
- PP 248
Mangal, Masa, "Modeling Fluctuations in Macromolecular Systems," 30 pp., Jun 1970
- PP 249
Trotz, Robert P., "The Estimation and Interpretation of General Selectivity Models," 37 pp., Jun 1970
- PP 250
Nunn, Walter R., "Position Finding with Prior Knowledge of Covariance Parameters," 8 pp., Jun 1970
- PP 251
Gleser, Kenneth S., "The d-Choice Secretary Problem," 32 pp., Jun 1970
- PP 252
Mangal, Masa and Guenbeck, David B., "Integration of a Bivariate Normal Over an Offset Circle," 14 pp., Jun 1970
- PP 253 - Classified
- PP 254
Thaler, R., "Discounting and Fixed Constraints: Why Discounting is Always Right," 10 pp., Aug 1970
- PP 255
Mangal, Masa S. and Thomas, James A., Jr., "Analytical Methods in Search Theory," 66 pp., Nov 1970
- PP 256
Goss, David V.; Hsu, In-Ching; Nunn, Walter R. and Pinto, David A., "A Class of Commutative Markov Matrices," 17 pp., Nov 1970 (To be submitted for publication in *Operations Research*)
- PP 257
Mangal, Masa S. and Goss, David K., "Detection Rate and Sweep Width in Visual Search," 14 pp., Nov 1970

END 1-80